

- STN SEARCH TRANSCRIPT - 10/800,328

MEDLINE ENABLEMENT

SEARCH FOR CLAIMED METHODS-OF-TREATMENT

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:ssspal623zct

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

***** Welcome to STN International *****

- NEWS 1 Web Page URLs for STN Seminar Schedule - N. America
 - NEWS 2 "Ask CAS" for self-help around the clock
 - NEWS 3 CA/Caplus pre-1967 chemical substance index entries enhanced with preparation role
 - NEWS 4 CA/Caplus patent kind codes updated
 - NEWS 5 MARPAT to CA/Caplus accession number crossover limit increased to 50,000
 - NEWS 6 DEC 18 MEDLINE updated in preparation for 2007 reload
 - NEWS 7 DEC 27 CA/Caplus enhanced with more pre-1907 records
 - NEWS 8 JAN 08 CHEMLIST enhanced with New Zealand Inventory of Chemicals
 - NEWS 9 JAN 16 CA/Caplus Company Name Thesaurus enhanced and reloaded
 - NEWS 10 JAN 16 IPC version 2007.01 thesaurus available on STN
 - NEWS 11 JAN 16 WPIDS/WPINDEX/WPIX enhanced with IPC 8 reclassification data
 - NEWS 12 JAN 22 CA/Caplus updated with revised CAS roles
 - NEWS 13 JAN 22 CA/Caplus enhanced with patent applications from India
 - NEWS 14 JAN 29 PHAR reloaded with new search and display fields
 - NEWS 15 JAN 29 CAS Registry Number crossover limit increased to 300,000 in multiple databases
 - NEWS 16 FEB 15 PATIDPASC enhanced with Drug Approval numbers
 - NEWS 17 FEB 15 RUSSJAPAT enhanced with pre-1994 records
 - NEWS 18 FEB 23 KOREAPAT enhanced with IPC 8 features and functionality
 - NEWS 19 FEB 26 MEDLINE reloaded with enhancements
 - NEWS 20 FEB 26 EMBASE enhanced with Clinical Trial Number field
 - NEWS 21 FEB 26 TOXCENTER enhanced with reloaded MEDLINE
 - NEWS 22 FEB 26 IFICDB/IEIPAT/IFIUDB reloaded with enhancements
 - NEWS 23 FEB 26 CAS Registry Number crossover limit increased from 10,000 to 300,000 in multiple databases
 - NEWS 24 MAR 15 WPIDS/WPIX enhanced with new FRAGHITSTR display format
 - NEWS 25 MAR 16 CASREACT coverage extended
 - NEWS 26 MAR 20 MARPAT now updated daily
 - NEWS 27 MAR 22 LWPI reloaded
 - NEWS 28 MAR 30 ROISCLOSURE reloaded with enhancements
 - NEWS 29 MAR 30 INPADOCDB will replace INPADOC on STN
 - NEWS 30 APR 02 JICST-EPLUS removed from database clusters and STN
- NEWS EXPRESS NOVEMBER 10 CURRENT WINDOWS VERSION IS V8.01C, CURRENT MACINTOSH VERSION IS V6.0c(ENG) AND V6.01c(JP), AND CURRENT DISCOVER FILE IS DATED 25 SEPTEMBER 2006.

- NEWS HOURS STN Operating Hours Plus Help Desk Availability
- NEWS LOGIN Welcome Banner and News Items
- NEWS IPC8 For general information regarding STN implementation of IPC 8
- NEWS X25 X.25 communication option no longer available

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN Customer agreement. Please note that this agreement limits use to scientific

***** STN Columbus *****

FILE 'HOME' ENTERED AT 12:29:16 ON 09 APR 2007

=> FILE MEDLINE

COST IN U.S. DOLLARS

FULL ESTIMATED COST

FILE 'MEDLINE' ENTERED AT 12:29:27 ON 09 APR 2007

FILE LAST UPDATED: 7 Apr 2007 (20070407/UP). FILE COVERS 1950 TO DATE.

This file contains CAS Registry Numbers for easy and accurate substance identification.

=> S BRIGHT

9184 BRIGHT

2 BRIGHTS

9185 BRIGHT

(BRIGHT OR BRIGHTS)

=> S L1 AND ANTIPSYCHOTICS

5343 ANTIPSYCHOTICS

1 L1 AND ANTIPSYCHOTICS

=> D

L2 ANSWER 1 OF 1 MEDLINE on STN

AN 2005667736 MEDLINE

DN PubMed ID: 16354123

TI Pharmacogenomics: a path to predictive medicine for schizophrenia.

AU Gupta Simone; Jain Sanjeev; Brahmachari Samir K; Kukreti Ritushree

CS Institute of Genomics and Integrative Biology (CSIR), Delhi University

Campus, Delhi 110007, India.

SO Pharmacogenomics, (2006 Jan) Vol. 7, No. 1, pp. 31-47. Ref: 159

Journal code: 100897350. ISSN: 1462-2416.

CY England; United Kingdom

DT Journal; Article; (JOURNAL ARTICLE)

(RESEARCH SUPPORT, NON-U.S. GOV'T)

General Review; (REVIEW)

LA English

FS Priority Journals

EM 200602

ED Entered STN: 20 Dec 2005

Last Updated on STN: 28 Feb 2006

Entered Medline: 27 Feb 2006

=> D ABS

L2 ANSWER 1 OF 1 MEDLINE on STN

AB A significant variability is observed among patients in response to antipsychotics, and is caused by a variety of factors. This review summarizes the available knowledge of associations between pharmacogenetics and drug response in schizophrenia. The multifactorial etiology of schizophrenia makes it a complex interaction of symptoms. Adopting a pharmacogenomics approach represents a unique opportunity for the prediction of response to antipsychotic drugs by investigating genes implicated with specific symptoms and side effects. A network model of the interaction/crosstalk between the neurotransmitter signaling systems

is presented to emphasize the importance of the genes associated with the molecular mechanisms of the disease and drug response. These genes may serve as potential susceptibility genes and drug targets for schizophrenia. The crucial point for the identification of a significant biologic marker(s) will include not only the experimental validation of the genes involved in the neurotransmitter signaling systems, but also the availability of large exactly comparable phenotyped patients samples. Coupling our knowledge of genetic polymorphisms with clinical response data promises a bright future for rapid advances in personalized medicine.

=> E BRIGHT GENE MICHAEL/AU 25
E1 17 BRIGHT G R/AU
E2 1 BRIGHT GAERTNER E/AU
E3 0 --> BRIGHT GENE MICHAEL/AU
E4 1 BRIGHT GEORGE M/AU
E5 7 BRIGHT H/AU
E6 57 BRIGHT H J/AU
E7 3 BRIGHT H L/AU
E8 2 BRIGHT HELEN/AU
E9 1 BRIGHT HOWARD/AU
E10 1 BRIGHT I/AU
E11 4 BRIGHT I B/AU
E12 1 BRIGHT ISAAC J/AU
E13 36 BRIGHT J/AU
E14 30 BRIGHT J E/AU
E15 1 BRIGHT J F/AU
E16 1 BRIGHT J I/AU
E17 20 BRIGHT J J/AU
E18 5 BRIGHT J L/AU
E19 33 BRIGHT J M/AU
E20 2 BRIGHT J N/AU
E21 7 BRIGHT J R/AU
E22 5 BRIGHT J S/AU
E23 1 BRIGHT JAMES E H/AU
E24 2 BRIGHT JANE H/AU
E25 8 BRIGHT JANICE M/AU

=> E WLODECKI BISHOP/AU 25
E1 1 WLODEWETZ V W/AU
E2 1 WLODECKA I/AU
E3 0 --> WLODECKI BISHOP/AU
E4 2 WLODECKI A/AU
E5 1 WLODEK AGATA/AU
E6 3 WLODEK B/AU
E7 23 WLODEK D/AU
E8 1 WLODEK DANUTA/AU
E9 1 WLODEK G/AU
E10 1 WLODEK G D/AU
E11 13 WLODEK G K/AU
E12 1 WLODEK J/AU
E13 23 WLODEK L/AU
E14 2 WLODEK L B/AU
E15 17 WLODEK LIDIA/AU
E16 1 WLODEK LIDIA B/AU
E17 2 WLODEK M/AU
E18 38 WLODEK M E/AU
E19 1 WLODEK MARY/AU
E20 6 WLODEK P J/AU
E21 1 WLODEK P J/AU
E22 1 WLODEK PRZENISLAW/AU
E23 2 WLODEK PRZENISLAW J/AU
E24 3 WLODEK S/AU
E25 6 WLODEK S T/AU

=> E BRODNEY MICHAEL A/AU 25
E1 1 BRODNEY K/AU
E2 3 BRODNEY M A/AU
E3 1 --> BRODNEY MICHAEL A/AU
E4 3 BRODNEY S/AU
E5 4 BRODNIAK W/AU
E6 2 BRODNIAK WLQDZIMIERZ A/AU
E7 3 BRODNIACK D/AU
E8 1 BRODNIACK D E/AU
E9 1 BRODNIACK DONALD E/AU
E10 1 BRODNIACK M S/AU
E11 2 BRODNIACK S/AU
E12 1 BRODNIACKA A/AU
E13 2 BRODNIACKA L/AU
E14 3 BRODNIACKI T/AU
E15 7 BRODNIACKI T C/AU
E16 2 BRODNIACKI THOMAS/AU
E17 8 BRODNIACKI THOMAS C/AU
E18 9 BRODNIEMICZ A/AU
E19 10 BRODNIEMICZ PROBA T/AU
E20 3 BRODNIEMICZ T/AU
E21 14 BRODNIEMICZ Z/AU
E22 2 BRODNIEMICZ V/AU
E23 1 BRODNIK ISABELLA/AU
E24 8 BRODNIK M/AU
E25 1 BRODNIK M M/AU

=> S E3
L3 1 "BRODNEY MICHAEL A"/AU

=> D IBIB ABS
L3 ANSWER 1 OF 1 MEDLINE on STN
ACCESSION NUMBER: 2004255135 MEDLINE
DOCUMENT NUMBER: Pubmed ID: 15153003
TITLE: A new strategy toward indole alkaloids involving an intramolecular cycloaddition/rearrangement cascade.
AUTHOR: Padwa Albert; Brodney Michael A; Lynch Stephen M; Rashatasakhon Paltoon; Wang Qiu; Zhang Hongjun
CORPORATE SOURCE: Department of Chemistry, Emory University, Atlanta, Georgia 30322, USA.. chemap@emory.edu
SOURCE: The Journal of organic chemistry, (2004 May 28) Vol. 69, No. 11, pp. 3735-45.
JOURNAL CODE: 2985193R. ISSN: 0022-3263.
PUB. COUNTRY: United States
DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)
LANGUAGE: English
FILE SEGMENT: Priority Journals
ENTRY MONTH: 200409
ENTRY DATE: Entered STN: 22 May 2004
Last Updated on STN: 21 Sep 2004
Entered Medline: 17 Sep 2004

AB The intramolecular Diels-Alder reaction between an amidofuran moiety tethered onto an indole component was examined as a strategy for the synthesis of Aspidosperma alkaloids. Furanyl carbamate 23 was acylated using the mixed anhydride 26 to provide amidofuran 22 in 68% yield. Further N-acylation of this indole furnished 27 in 88% yield. Cyclization precursors were prepared by removing the carbamate moiety followed by N-alkylation with the appropriate alkyl halides. Large substituent groups on the amido nitrogen atom causes the reactive s-trans conformation of the amidofuran to be more highly populated, thereby facilitating the Diels-Alder cycloaddition. The reaction requires the presence of an electron-withdrawing substituent on the indole nitrogen in order for the

cycloaddition to proceed. Treatment of N-allyl-bromenamide 48 with n-Bu(3)ShH/AIBN preferentially led to the 6-endo trig cyclization product 50, with the best yield (91%) being obtained under high dilution conditions. The initially generated cyclohexenyl radical derived from 48 produces the pentacyclic heterocycle 50 by either a direct 6-endo trig cyclization or, alternatively, by a vinyl radical rearrangement pathway.

=> S ANTAGONISTS AND (5HT1B OR 5HT2A OR D2)
487456 ANTAGONISTS
137 5HT1B
179 5HT2A
26201 D2

L4 6832 ANTAGONISTS AND (5HT1B OR 5HT2A OR D2)

=> S L4 AND 2003/PY
569314 2003/PY
(20030000-20039999/PY)
L5 349 L4 AND 2003/PY

=> S L5 AND REVIEW
475081 REVIEW
59625 REVIEWS
520853 REVIEW
(REVIEW OR REVIEWS)

L6 7 L5 AND REVIEW

=> D 1-7 1B1B ABS

L6 ANSWER 1 OF 7 MEDLINE on STN
ACCESSION NUMBER: 2004019152 MEDLINE
DOCUMENT NUMBER: Pubmed ID: 14715440
TITLE: Novel mechanisms and approaches in the study of neurodegeneration and neuroprotection. a review.
AUTHOR: Kostrewa Richard M; Segura-Aguilar Juan
CORPORATE SOURCE: Department of Pharmacology, Quillen College of Medicine, East Tennessee State University, Johnson City, TN 37614, USA.. kostrew@etsu.edu

CONTRACT NUMBER: NS 39272 (NINDS)
SOURCE: Neurotoxicity research, (2003) Vol. 5, No. 6, pp. 375-83. Ref: 93
Journal code: 100929017. ISSN: 1029-8428.

PUB. COUNTRY: United States
DOCUMENT TYPE: Journal: Article; (JOURNAL ARTICLE)
(RESEARCH SUPPORT, NON-U.S. GOV'T)
(RESEARCH SUPPORT, U.S. GOV'T, P.H.S.)
General Review; (REVIEW)

LANGUAGE: English
FILE SEGMENT: Priority Journals
ENTRY MONTH: 200402
ENTRY DATE: Entered STN: 13 Jan 2004
Last Updated on STN: 3 Feb 2004

Entered Medline: 2 Feb 2004

AB Cellular mechanisms involved in neurodegeneration and neuroprotection are continuing to be explored, and this paper focuses on some novel discoveries that give further insight into these processes. Oligodendrocytes and activated astroglia are likely generators of the pro-inflammatory cytokines, such as the tumor necrosis factor family and interleukin family, and these glial support cells express adhesion receptors (e.g., VCAM) and release intercellular adhesion molecules (ICAM) that have a major role in neuronal apoptosis. Even brief exposure to some substances, in ontogeny and sometimes in adulthood, can have lasting effects on behaviors because of their prominent toxicity (e.g., NMDA receptor antagonists) or because they sensitize receptors (e.g., dopamine D2 agonists), possibly permanently, and thereby alter

behavior for the lifespan. Cell cycle genes which may be derived from microglia, are the most-recent entry into the neuroprotection schema. Neuroprotection afforded by some common substances (e.g., melatonin) and uncommon substances (e.g., nicotine, green tea polyphenol (-)-epigallocatechin-3-gallate (EGCG), trolox), ordinarily thought to be simple radical scavengers, now are thought to invoke previously unsuspected cellular mechanisms in the process of neuroprotection.

Although Alzheimer's disease (AD) has features of a continuous spectrum of neural and functional decline, in vivo PET imaging and functional magnetic resonance imaging, indicate that AD can be staged into an early phase treatable by inhibitors of beta and gamma secretase, and a late phase which may be more amenable to treatment by drugs that prevent or reverse tau phosphorylation. Neural transplantation, thought to be the last hope for neurally injured patients (e.g., Parkinsonians), may be displaced by non-neural tissue transplants (e.g., human umbilical cord blood; Sertoli cells) which seem to provide similar neurotrophic support and improved behavior - without posing the major ethical dilemma of removing tissue from aborted fetuses. The objective of this paper is to invite added research into the newly discovered (or postulated) novel mechanisms; and to stimulate discovery of additional mechanisms attending neurodegeneration and neuroprotection.

L6 ANSWER 2 OF 7 MEDLINE on STN
ACCESSION NUMBER: 2003582573 MEDLINE
DOCUMENT NUMBER: Pubmed ID: 14663001

TITLE: Adenosine-dopamine interactions: development of a concept and some comments on therapeutic possibilities.

AUTHOR: Fredholm Bertil B; Svenningsson Per
CORPORATE SOURCE: Department of Physiology and Pharmacology, Karolinska Institutet, Stockholm, Sweden.. Bertil.Fredholm@yfa.ki.se
SOURCE: Neurology, (2003 Dec 9) Vol. 61, No. 11 Suppl 6, pp. S5-9. Ref: 63
Journal code: 0401060. E-ISSN: 1526-632X.

PUB. COUNTRY: United States

DOCUMENT TYPE: Journal: Article; (JOURNAL ARTICLE)

LANGUAGE: English

FILE SEGMENT: Abridged Index Medicus Journals; Priority Journals

ENTRY MONTH: 200402

ENTRY DATE: Entered STN: 16 Dec 2003

Last Updated on STN: 12 Feb 2004

Entered Medline: 11 Feb 2004

AB This brief review presents a personal perspective on the historical development of the current knowledge about the biologically important concept of functional antagonism between adenosine A2A and dopamine D2 receptors in caudate-putamen, accumbens, and tuberculum olfactorium. In the 1970s, studies of dopamine actions suggested an unexpected role of adenosine. Developments during the next decade substantiated this finding and demonstrated that a subform of adenosine A2 receptors was enriched in the basal ganglia. Cloning of adenosine receptors provided better tools for cellular localization and showed that A2A receptors are closely associated with D2 receptors. Distinct functional interactions at several levels were discovered, and there is now strong evidence that A2A receptors are tonically active and modified by dopamine acting at D2 receptors. Development of selective antagonists and knockout mice have highlighted the potential usefulness of A2A antagonists in decreasing symptoms and progression of Parkinson's disease-something that has also been vindicated by careful epidemiologic studies. There are issues of efficacy and potential side effects that need to be resolved, but the future looks bright.

L6 ANSWER 3 OF 7 MEDLINE on STN
ACCESSION NUMBER: 2003565927 MEDLINE
DOCUMENT NUMBER: Pubmed ID: 14647530

TITLE: Neuropharmacological profile of an atypical antipsychotic, NRA0562.

AUTHOR: Hirota Shino; Kawashima Naoya; Chaki Shigeyuki; Okuyama Shigeru

CORPORATE SOURCE: Psychiatric Diseases and Pain Research, Medicinal Pharmacology Laboratory, Medicinal Research Laboratories, Taisho Pharmaceutical Co., Ltd., 1-403 Yoshino-cho, Kira-ku, Saitama, Saitama 331-9530, Japan.. shino.hirota@rd.taisho.co.jp

SOURCE: CNS drug reviews, (2003 Winter) Vol. 9, No. 4, pp. 375-88. Ref: 70

PUB. COUNTRY: United States

DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)

LANGUAGE: English

FILE SEGMENT: Priority Journals

ENTRY MONTH: 200401

ENTRY DATE: Entered STN: 16 Dec 2003
Last Updated on STN: 21 Jan 2004
Entered Medline: 20 Jan 2004

AB Schizophrenia is a serious and disabling psychiatric disorder affecting approximately 1% of the world's population. A new generation of atypical antipsychotics has been introduced over the past decade. These atypical antipsychotics have comparable or greater efficacy than traditional antipsychotics in the treatment of the psychotic symptoms of schizophrenia and a much improved neurologic side effect profile. This paper reviews the pharmacological efficacy and safety of a potential atypical antipsychotic, NRA0562. NRA0562 has a high affinity for dopamine D1, D2L, D4.2, 5-HT2A receptors as well as alpha1-adrenoceptors, and has a moderate affinity for H1 receptors. NRA0562 strongly binds to 5-HT2A receptors and alpha1-adrenoceptors in the frontal cortex, its binding to striatal D2 receptors is weaker, similar to that of clozapine. NRA0562 displayed potent antipsychotic activities in animal models of schizophrenia, such as methamphetamine (MAP)-induced hyperactivity, apomorphine-induced disruption of pre-pulse inhibition and conditioned avoidance test. NRA0562 is more potent in reversing the inhibitory effects of MAP at A10 than at A9 dopamine neurons. It increased Fos-like immunoreactivity in the nucleus accumbens more effectively than in the dorsolateral striatum, indicating that NRA0562 has the profile of an atypical antipsychotic. In vivo assays for extrapyramidal side effect liability showed that NRA0562 has a low rate of neurological side effects. Thus, NRA0562 may have unique antipsychotic activity with a lower propensity for extrapyramidal side effects.

L6 ANSWER 4 OF 7 MEDLINE on STN

ACCESSION NUMBER: 2003533618 MEDLINE

DOCUMENT NUMBER: PubMed ID: 14612145

TITLE: Rapid regulation of dopamine transporter function by substrates, blockers and presynaptic receptor ligands.

AUTHOR: Gulley Joshua M; Zahnliser Nancy R

CORPORATE SOURCE: Department of Pharmacology and Neuroscience Program, University of Colorado Health Sciences Center, Campus Box C-236, 4200 E Ninth Avenue, Denver, CO 80262, USA.. joshua.gulley@uchsc.edu

CONTRACT NUMBER: DA 04216 (NIDA)

SOURCE: DA 15050 (NIDA)
European Journal of pharmacology, (2003 Oct 31) Vol. 479, No. 1-3, pp. 139-52. Ref: 140
Journal code: 1254354. ISSN: 0014-2999.

PUB. COUNTRY: Netherlands

DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)

LANGUAGE: (RESEARCH SUPPORT, U.S. GOV'T, P.H.S.)
General Review; (REVIEW)
English

FILE SEGMENT: Priority Journals

ENTRY MONTH: 200407

ENTRY DATE: Entered STN: 13 Nov 2003
Last Updated on STN: 21 Jul 2004
Entered Medline: 20 Jul 2004

AB The extracellular actions of dopamine are terminated primarily through its binding to dopamine transporters and translocation back into dopamine neurons. The transporter thereby serves as an optimal target to regulate dopamine neurotransmission. Although acute pharmacological blockade of dopamine transporters is known to reversibly inhibit transporter function by preventing the binding of its endogenous substrate dopamine, it recently has become clear that dopamine transporter substrates, such as amphetamines, and blockers, such as cocaine, also have the ability to rapidly and persistently regulate transporter function after their direct pharmacological effect has subsided. Presynaptic receptor ligands can also regulate dopamine transporter function. This has been investigated most extensively for dopamine D2 receptors, but there is also evidence for regulation by gamma-aminobutyric acid (GABA) GABAB receptors, metabotropic glutamate, nicotinic acetylcholine, serotonin, sigma2- and kappa-opioid receptors. The focus of this review is the rapid, typically reversible, regulation of dopamine transporter velocity by substrates, blockers and presynaptic receptor ligands. The research discussed here suggests that a common mechanism through which these different classes of compounds regulate transporter activity is by altering the cell surface expression of dopamine transporters.

L6 ANSWER 5 OF 7 MEDLINE on STN

ACCESSION NUMBER: 2003364198 MEDLINE

DOCUMENT NUMBER: PubMed ID: 12895600

TITLE: The second PGD(2) receptor CRTH2: structure, properties, and functions in leukocytes.

AUTHOR: Nagata Kinya; Hirai Hiroyuki

CORPORATE SOURCE: R&D Centre, Bio Medical Laboratories, Inc, 1361-1 Matoba, Kawagoe, Saitama 350-1101, Japan.. nagata@alk.co.jp

SOURCE: Prostaglandins, leukotrienes, and essential fatty acids, (2003 Aug-Sep) Vol. 69, No. 2-3, pp. 169-77. Ref: 76

PUB. COUNTRY: Scotland; United Kingdom

DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)

LANGUAGE: English

FILE SEGMENT: Priority Journals

ENTRY MONTH: 200404

ENTRY DATE: Entered STN: 5 Aug 2003
Last Updated on STN: 17 Apr 2004
Entered Medline: 16 Apr 2004

AB Prostaglandin (PG) D(2) plays a broad range of physiological and pathophysiological functions. Until just a few years ago, it was thought that most of the biological actions of PGD(2) are mediated via the classical PGD(2) receptor DP. Recently, we identified a second PGD(2) receptor, chemoattractant receptor-homologous molecule expressed on T helper (Th)2 cells (CRTH2), with different functions relative to DP. Here, we review the recent findings on the structure, tissue distribution, ligand selectivity, signalling pathways, and functions in leukocytes of this receptor. The data suggest that the PGD(2)/CRTH2 system play important roles in allergic inflammation through its stimulatory effects on Th2 cells, eosinophils, and basophils.

L6 ANSWER 6 OF 7 MEDLINE on STN

ACCESSION NUMBER: 2003288164 MEDLINE

DOCUMENT NUMBER: PubMed ID: 12814658

TITLE: Targeting striatal cholinergic interneurons in Parkinson's disease: focus on metabotropic glutamate receptors.

AUTHOR: Pisani A; Bonsi P; Centonze D; Cubellini P; Bernardi G; Calabresi P
CORPORATE SOURCE: Clinica Neurologica, Dipartimento di Neuroscienze, Università di Roma Tor Vergata, Rome, Italy..
 pisani@uniroma2.it
SOURCE: Neuropsychopharmacology, (2003 Jul) Vol. 45, No. 1, pp. 45-56. Ref: 95
PUB. COUNTRY: Journal code: 0236217, ISSN: 0028-3908.
DOCUMENT TYPE: Journal: United Kingdom
 (RESEARCH SUPPORT, NON-U.S. GOV'T)
LANGUAGE: English
ENTRY MONTH: Priority Journals
ENTRY DATE: 200308
 Entered STN: 20 Jun 2003
 Last Updated on STN: 12 Aug 2003
 Entered Medline: 11 Aug 2003

AB In the early sixties, anticholinergic drugs were introduced in the pharmacological treatment of Parkinson's disease (PD). The rationale behind their utilisation in the treatment of the disease was based on the evidence of an imbalance between the dopaminergic inputs and the intrinsic cholinergic innervation within the striatum. Metabotropic glutamate (mGlu) receptors have been shown to play a key role in striatal function both in physiological conditions and in experimental models of diseases affecting this brain area. Indeed, compelling electrophysiological and morphological evidence shows that mGlu receptors are highly expressed at cellular level and exert a profound modulatory role on cholinergic interneurons excitability. This review will provide a brief survey of studies on the localization and function of mGlu receptors in cholinergic interneurons. The potential relevance of these findings in the control of motor function and in the treatment of PD will be discussed.

L6 ANSWER 7 OF 7
ACCESSION NUMBER: MEDLINE on STN
DOCUMENT NUMBER: 2003275541
TITLE: PubMed ID: 12801600
 Neurotensin: dual roles in psychostimulant and antipsychotic drug responses.
AUTHOR: Dobner Paul R; Deutch Ariel Y; Fadel Jim
CORPORATE SOURCE: Department of Molecular Genetics and Microbiology, Program in Neuroscience, University of Massachusetts Medical School, 55 Lake Ave. North, Worcester 01655, USA..
 paul.dobner@umassmed.edu
CONTRACT NUMBER: HL-33307 (NHLBI)
 MH-45124 (NIMH)
 NS-44282 (NINDS)
SOURCE: Life sciences, (2003 Jun 27) Vol. 73, No. 6, pp. 801-11. Ref: 82
PUB. COUNTRY: Journal code: 0375521, ISSN: 0024-3205.
DOCUMENT TYPE: Journal: United Kingdom
 (RESEARCH SUPPORT, NON-U.S. GOV'T)
 General Review: (REVIEW)
LANGUAGE: English
FILE SEGMENT: Priority Journals
ENTRY MONTH: 200307
ENTRY DATE: Entered STN: 13 Jun 2003
 Last Updated on STN: 18 Jul 2003
AB Central administration of neurotensin (NT) results in a variety of neurobehavioral effects which, depending upon the administration site, resemble the effects of antipsychotic drugs (APDs) and psychostimulants.

All clinically effective APDs exhibit significant affinities for dopamine D(2) receptors, supporting the hypothesis that an increase in dopaminergic tone contributes to schizophrenic symptoms. Psychostimulants increase extracellular dopamine (DA) levels and chronics administration can produce psychotic symptoms over time. APDs and psychostimulants induce Fos and NT expression in distinct striatal subregions, suggesting that changes in gene expression underlie some of their effects. To gain insight into the functions of NT, we analyzed APD and psychostimulant induction of Fos in NT knockout mice and rats pretreated with the NT antagonist SR 48692. In both NT knockout mice and rats pretreated with SR 48692, haloperidol-induced Fos expression was markedly attenuated in the dorsolateral striatum; amphetamine-induced Fos expression was reduced in the medial striatum. These results indicate that NT is required for the activation of specific subpopulations of striatal neurons in distinct striatal subregions in response to both APDs and psychostimulants. This review integrates these new findings with previous evidence implicating NT in both APD and psychostimulant responses.

=> FILE REG COST IN U.S. DOLLARS SINCE FILE ENTRY TOTAL
 FULL ESTIMATED COST 15.24 15.45

FILE 'REGISTRY' ENTERED AT 12:49:42 ON 09 APR 2007
 USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
 PLEASE SEE "HELP USAGETERMS" FOR DETAILS
 COPYRIGHT (C) 2007 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file provided by InfoChem.

STRUCTURE FILE UPDATES: 8 APR 2007 HIGHEST RN 929518-97-8
 DICTIONARY FILE UPDATES: 8 APR 2007 HIGHEST RN 929518-97-8

New CAS Information Use Policies, enter HELP USAGETERMS for details.

TSCA INFORMATION NOW CURRENT THROUGH December 2, 2006
 Please note that search-term pricing does apply when conducting SmartSELECT searches.

REGISTRY includes numerically searchable data for experimental and predicted properties as well as tags indicating availability of experimental property data in the original document. For information on property searching in REGISTRY, refer to:

<http://www.cas.org/ONLINE/UG/regprops.html>

=> S NRA0562/CN
 L7 0 NRA0562/CN

=> S NRA0562
 L8 0 NRA0562

=> SEL 3
 L8 HAS NO ANSWERS
 An L-number has no answers for one of five reasons:
 1. It is a query that has not been searched, or
 2. It is the result of a search with zero answers, or
 3. It is an intermediate result of the ACTIVATE command, or
 4. It is an intermediate result in SEARCH STEPS, or
 5. It is an L-number created by the RUN command

```

=> SEL L6 3
NO ANSWERS SELECTED. THE ANSWER SET WAS CREATED IN FILE 'MEDLINE'.
USE THE FILE COMMAND TO CHANGE TO THE CORRECT FILE.
An answer set can be processed to create terms only
in the same file in which it was created.

=> FILE MEDLINE
COST IN U.S. DOLLARS SINCE FILE TOTAL
ENTRY 10.80 SESSION 26.25
FULL ESTIMATED COST

FILE 'MEDLINE' ENTERED AT 12:50:36 ON 09 APR 2007
FILE LAST UPDATED: 7 Apr 2007 (20070407/UP). FILE COVERS 1950 TO DATE.

This file contains CAS Registry Numbers for easy and accurate
substance identification.

=> SEL L6 3
E1 THROUGH E7 ASSIGNED

=> FIL STNGUIDE
COST IN U.S. DOLLARS SINCE FILE TOTAL
ENTRY 0.39 SESSION 26.64
FULL ESTIMATED COST

FILE 'STNGUIDE' ENTERED AT 12:51:24 ON 09 APR 2007
USE IS SUBJECT TO THE TERMS OF YOUR CUSTOMER AGREEMENT
COPYRIGHT (C) 2007 AMERICAN CHEMICAL SOCIETY, JAPAN SCIENCE
AND TECHNOLOGY CORPORATION, AND FACHINFORMATIONSZENTRUM KARLSRUHE

FILE CONTAINS CURRENT INFORMATION.
LAST RELOADED: Apr 6, 2007 (20070406/UP).

=> FILE MEDLINE
COST IN U.S. DOLLARS SINCE FILE TOTAL
ENTRY 0.06 SESSION 26.70
FULL ESTIMATED COST

FILE 'MEDLINE' ENTERED AT 12:52:14 ON 09 APR 2007
FILE LAST UPDATED: 7 Apr 2007 (20070407/UP). FILE COVERS 1950 TO DATE.

This file contains CAS Registry Numbers for easy and accurate
substance identification.

=> D E1-E7
'E1-E7' IS NOT A VALID ACCESSION NUMBER

The number entered is not a valid accession number in this file.
Enter "HELP ACCESSION" at an arrow prompt (=>) for a list of valid
accession number formats in the current file.

=> S 2003565927/AN
L9 1 2003565927/AN

=> D
L9 ANSWER 1 OF 1 MEDLINE on STN
AN 2003565927 MEDLINE
DN PubMed ID: 14647530
TI Neuropharmacological profile of an atypical antipsychotic, NRA0562.
AU Hirota Shihou; Kawashima Naoya; Chaki Shigeyuki; Okuyama Shigeru
CS Psychiatric Diseases and Pain Research, Medicinal Pharmacology Laboratory,
Medicinal Research Laboratories, Taisho Pharmaceutical Co., Ltd., 1-403

```

```

Yoshino-cho, Kita-ku, Saitama, Saitama 331-9530, Japan..
shino.hirota@po.rd.taisho.co.jp
CNS drug reviews, (2003 Winter) Vol. 9, No. 4, pp. 375-88. Ref: 70
Journal code: 9514898. ISSN: 1080-563X.
CY United States
DT Journal; Article; (JOURNAL ARTICLE)
LA English
FS Priority Journals
EM 200401
ED Entered STN: 16 Dec 2003
Last Updated on STN: 21 Jan 2004
Entered Medline: 20 Jan 2004

=> SEL RN
E8 THROUGH E8 ASSIGNED

=> D E8
'E8' IS NOT A VALID FORMAT FOR FILE 'MEDLINE'

The following are valid formats:
The default display format is BIB.

ABS ---- AB
ALL ---- AN, DN, TI, AU, CS, NC, SO, CM, CY, DT, LA, FS, NCT, OS,
EM, ED, AB, ST, CT, NA, RN, CN, GEN
BIB ---- AN, DN, TI, AU, CS, NC, SO, CY, DT, LA, FS, NCT, OS, EM, ED
CBIB ---- AN, DN, TI, AU, CS, NC, SO, CY, DT, LA, FS, NCT, OS, EM, ED
DALL ---- ALL, delimited for post processing
IABS ---- ABS, with a text label
IALL ---- ALL, indented with text labels
IBIB ---- BIB, indented with text labels
IND ---- ST, CT, NA, RN, CN, GEN
TRIAL -- TI, ST, CT, NA, RN, CN, GEN
(SAM, TRI, FREE)
HIT ---- All fields containing hit terms
HITIND - IND
KWIC ---- All hit terms plus 20 words on either side
OCC ---- List of display fields containing hit terms

Hit terms will be highlighted in all available fields except CM and
PY.

To display a particular field or fields, enter the display field
codes. For a list of display field codes, enter 'HELP DFIELDS'
at an arrow prompt (=>). Examples of formats include: 'BIB',
'AB', 'SO,ST'. You may specify the format fields in any order,
and the information will be displayed in the same order as the
format specification.

The same formats (except for HIT, HITIND, KWIC, and OCC) may be
used with the DISPLAY ACC command to display the record for a
specified Accession Number.
ENTER DISPLAY FORMAT (BIB):END

=> FILE REG
COST IN U.S. DOLLARS SINCE FILE TOTAL
ENTRY 0.56 SESSION 27.26
FULL ESTIMATED COST


FILE 'REGISTRY' ENTERED AT 12:53:04 ON 09 APR 2007
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

```

COPYRIGHT (C) 2007 American Chemical Society (ACS)
Property values tagged with IC are from the ZIC/VINITI data file
provided by InfoChem.
STRUCTURE FILE UPDATES: 8 APR 2007 HIGHEST RN 929518-97-8
DICTIONARY FILE UPDATES: 8 APR 2007 HIGHEST RN 929518-97-8
New CAS Information Use Policies, enter HELP USAGETERMS for details.
TSCA INFORMATION NOW CURRENT THROUGH December 2, 2006
Please note that search-term pricing does apply when
conducting SmartSELECT searches.
REGISTRY includes numerically searchable data for experimental and
predicted properties as well as tags indicating availability of
experimental property data in the original document. For information
on property searching in REGISTRY, refer to:
<http://www.cas.org/ONLINE/UG/regprops.html>

=> S E8
L10 1 537-46-2/BI
(537-46-2/RN)
=> D
L10 ANSWER 1 OF 1 REGISTRY COPYRIGHT 2007 ACS on STN
RN 537-46-2 REGISTRY
ED Entered STN: 16 Nov 1984
CN Benzenethanamine, N,α-dimethyl-, (αS)- (CA INDEX NAME)
OTHER CA INDEX NAMES:
CN Benzenethanamine, N,α-dimethyl-, (S)-
CN Phenethylamine, N,α-dimethyl-, (S)-(+)- (8CI)
OTHER NAMES:
CN (+)-(S)-Deoxyephedrine
CN (+)-2-(N-Methylamino)-1-phenylpropane
CN (+)-Methamphetamine
CN (+)-Methylamphetamine
CN (+)-N,α-Dimethyl-β-phenylethylamine
CN (+)-N-Methylamphetamine
CN (S)-(+)-Deoxyephedrine
CN (S)-(+)-Methamphetamine
CN (S)-Methamphetamine
CN 2S-(+)-Methamphetamine
CN Corvatin
CN d-(S)-Methamphetamine
CN d-Deoxyephedrine
CN d-Deoxyephedrine
CN d-Methamphetamine
CN d-Methylamphetamine
CN d-N,α-Dimethylphenethylamine
CN d-N-Methylamphetamine
CN d-Phenylisopropylmethylamine
CN L-Methamphetamine
CN Metamphetamine
CN Metamphetamine
CN Methyl-β-phenylisopropylamine
CN Methylamphetamine
CN N-Methyl-1-phenyl-2-propanamine
CN N-Methylamphetamine
CN Norodin

CN NSC 25115
FS STEREOSEARCH
DR 139-47-9, 1690-86-4, 14611-50-8, 45952-89-4
MF C10 H15 N
CI COM
LC STN Files: ADISNEWS, AGRICOLA, ANABSTR, BEILSTEIN*, BIOSIS, BIOTECHNO,
CA, CAOLD, CAPLUS, CASREACT, CBNB, CHEMCATS, CHEMLIST, CIN, CSCHEM,
CSNB, DDFU, DRUGU, EMBASE, HSDB*, IFICDB, IFIPAT, IFIUDE, IPA, MEDLINE,
MRCK*, PIRA, PROMT, PS, RTECS*, SCISEARCH, SPECINFO, SYNTHLINE,
TOXENTER, USAN, USPAT2, USPATFULL
(*File contains numerically searchable property data)
Other Sources: EINECS**, WHO
(*Enter CHEMLIST File for up-to-date regulatory information)
Absolute stereochemistry. Rotation (+).



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

4580 REFERENCES IN FILE CA (1907 TO DATE)
100 REFERENCES TO NON-SPECIFIC DERIVATIVES IN FILE CA
4597 REFERENCES IN FILE CAPLUS (1907 TO DATE)
19 REFERENCES IN FILE CAOLD (PRIOR TO 1967)

=> LOG HOLD
COST IN U.S. DOLLARS
FULL ESTIMATED COST
SINCE FILE ENTRY 3.75
TOTAL SESSION 31.01

SESSION WILL BE HELD FOR 120 MINUTES
STN INTERNATIONAL SESSION SUSPENDED AT 12:55:36 ON 09 APR 2007
Connecting via Winsock to STN

Welcome to STN International! Enter x:x
LOGINID:sssptal623zct

PASSWORD:
***** * RECONNECTED TO STN INTERNATIONAL * * * * *
SESSION RESUMED IN FILE 'REGISTRY' AT 13:04:58 ON 09 APR 2007
FILE 'REGISTRY' ENTERED AT 13:04:58 ON 09 APR 2007
COPYRIGHT (C) 2007 American Chemical Society (ACS)

COST IN U.S. DOLLARS
FULL ESTIMATED COST
SINCE FILE ENTRY 3.75
TOTAL SESSION 31.01

=> FILE 2002/PY
'2002/PY' IS NOT A VALID FILE NAME
SESSION CONTINUES IN FILE 'REGISTRY'
Enter "HELP FILE NAMES" at an arrow prompt (=>) for a list of files
that are available. If you have requested multiple files, you can
specify a corrected file name or you can enter "IGNORE" to continue
accessing the remaining file names entered.

=> FILE MEDLINE
COST IN U.S. DOLLARS
FULL ESTIMATED COST

SINCE FILE ENTRY 4.20
TOTAL SESSION 31.46

FILE 'MEDLINE' ENTERED AT 13:05:22 ON 09 APR 2007

FILE LAST UPDATED: 7 Apr 2007 (20070407/UP). FILE COVERS 1950 TO DATE.

This file contains CAS Registry Numbers for easy and accurate substance identification.

=> S 2002/PY
L11 542785 2002/PY
(20020000-20029999/PY)

=> S L11 AND REVIEW
475081 REVIEW
59625 REVIEWS
520853 REVIEW
(REVIEW OR REVIEWS)

L12 30323 L11 AND REVIEW
116 BENZISOXAZOLE
20 BENZISOXAZOLES
130 BENZISOXAZOLE
(BENZISOXAZOLE OR BENZISOXAZOLES)

L13 0 L12 AND BENZISOXAZOLE

=> S BENZISOXAZOLE AND 2003/PY
116 BENZISOXAZOLE
120 BENZISOXAZOLES
130 BENZISOXAZOLE
(BENZISOXAZOLE OR BENZISOXAZOLES)

569314 2003/PY
(20030000-20039999/PY)

L14 5 BENZISOXAZOLE AND 2003/PY

=> D 1-5

L14 ANSWER 1 OF 5 MEDLINE on STN
AN 2003569076 MEDLINE
DN PubMed ID: 14640551
TI 1,2-benzisoxazole phosphorodiamidates as novel anticancer
prodrugs requiring bio-reductive activation.
AU Jain Monish; Kwon Chul-Hoon
CS Department of Pharmaceutical Sciences, College of Pharmacy and Allied
Health Professions, St John's University, Jamaica, New York 11439, USA.
SO Journal of medicinal chemistry, (2003 Dec 4) Vol. 46, No. 25,
pp. 5428-36.
Journal code: 9716531. ISSN: 0022-2623.

CY United States
DT (IN VITRO)
LA English
FS Priority Journals
EM 200401
ED Entered STN: 16 Dec 2003
Last Updated on STN: 17 Jan 2004
Entered Medline: 16 Jan 2004

L14 ANSWER 2 OF 5 MEDLINE on STN
AN 2003405077 MEDLINE

DN PubMed ID: 12944663
TI 2-(2,1-benzoxazol-3-yl)-3,5-dimethoxyphenol and 3-phenyl-2,1-benzoxazole.
AU Howie R Alan; Jabbar Abdul; Lewis John R; Nizami Shaikh S; Ritchie Craig F
CS Department of Chemistry, University of Aberdeen, Meston Walk, Aberdeen
AB24 3UE, Scotland.. r.a.howie@abdn.ac.uk
SO Acta crystallographica. Section C, Crystal structure communications,
(2003 Sep) Vol. 59, No. Pt 9, pp. o516-9. Electronic Publication:
2003-08-09.
Journal code: 8305826. ISSN: 0108-2701.

CY Denmark
DT Journal; Article; (JOURNAL ARTICLE)
LA English
FS NONMEDLINE; PUBMED-NOT-MEDLINE
EM 200402
ED Entered STN: 29 Aug 2003
Last Updated on STN: 2 Mar 2004
Entered Medline: 26 Feb 2004

L14 ANSWER 3 OF 5 MEDLINE on STN
AN 200333232 MEDLINE
DN PubMed ID: 12867488
TI Substituent effect on the reductive N-dearylation of 3-(indol-1-yl)-1,2-
benzisoxazoles by rat liver microsomes.
AU Tschirret-Guth Richard A; Wood Harold B
CS Department of Drug Metabolism, Merck Research Laboratories, Rahway, NJ
07065, USA.. richard.tschirretguth@merck.com
SO Drug metabolism and disposition: the biological fate of chemicals,
(2003 Aug) Vol. 31, No. 8, pp. 999-1004.
Journal code: 9421550. ISSN: 0090-9556.

CY United States
DT Journal; Article; (JOURNAL ARTICLE)
LA English
FS Priority Journals
EM 200406
ED Entered STN: 18 Jul 2003
Last Updated on STN: 18 Jun 2004
Entered Medline: 17 Jun 2004

L14 ANSWER 4 OF 5 MEDLINE on STN
AN 2003141915 MEDLINE
DN PubMed ID: 12657263
TI Phenylacetic acid derivatives as hPPAR agonists.
AU Santini Conrad; Berger Gregory D; Han Wei; Mosley Ralph; MacNaul Karen;
Berger Joel; Doeber Thomas; Wu Margaret; Moller David E; Tolman Richard
L; Sahoo Soumya P
CS Department of Basic Chemistry, Merck Research Laboratories, Rahway, NJ
07065, USA.. conrad_santini@merck.com
SO Bioorganic & medicinal chemistry letters, (2003 Apr 7) Vol. 13,
No. 7, pp. 1277-80.
Journal code: 9107377. ISSN: 0960-894X.

CY England; United Kingdom
DT Journal; Article; (JOURNAL ARTICLE)
LA English
FS Priority Journals
EM 200311
ED Entered STN: 27 Mar 2003
Last Updated on STN: 4 Nov 2003
Entered Medline: 3 Nov 2003

L14 ANSWER 5 OF 5 MEDLINE on STN
AN 2003073789 MEDLINE
DN PubMed ID: 12583727
TI Isoxazole --> benzisoxazole rearrangement promoted cascade
reactions affording stereodefined polycycles.
AU Bode Jeffrey W; Uesaka Hidehiro; Suzuki Keisuke

CS Department of Chemistry, Tokyo Institute of Technology, and CREST, Japan
Science and Technology (JST) Corporation, O-okayama, Meguro-ku, Tokyo
152-8531, Japan. (2003 Feb 20) Vol. 5, No. 4, pp. 395-8.
SO Organic letters. ISSN: 100890393. ISSN: 1523-7060.
CY United States
DT Journal; Article; (JOURNAL ARTICLE)
LA English
FS NONMEDLINE; PUBMED-NOT-MEDLINE
EM 200307
ED Entered STN: 14 Feb 2003
Last Updated on STN: 17 Jul 2003
Entered Medline: 16 Jul 2003

=> S BENZISOXAZOLE AND (D2 OR 5HT1B OR 5HT2B OR 5HT-1A OR 5HT-2B)

116 BENZISOXAZOLE
20 BENZISOXAZOLES
130 BENZISOXAZOLE
(BENZISOXAZOLE OR BENZISOXAZOLES)

26201 D2
137 5HT1B
11 5HT2B
2604 5HT
19985 1A
104 5HT-1A
(5HT(W)1A)
2604 5HT
17924 2B
3 5HT-2B
(5HT(W)2B)

L15 14 BENZISOXAZOLE AND (D2 OR 5HT1B OR 5HT2B OR 5HT-1A OR 5HT-2B)

=> D 1-14 IBIB ABS

L15 ANSWER 1 OF 14 MEDLINE on STN
ACCESSION NUMBER: 2005342126
DOCUMENT NUMBER: Pubmed ID: 15992090
TITLE: Risperidone (Risperdal): clinical experience with a new antipsychosis drug.
AUTHOR: Keks N A; Culhane C
CORPORATE SOURCE: Monash University, Mental Health Research Institute of Victoria, Alfred Hospital, Prahran 3181, Australia..
N.Keks@alfred.org.au
SOURCE: Expert opinion on investigational drugs, (1999 Apr) Vol. 8, No. 4, pp. 443-52.
Journal code: 9434197. E-ISSN: 1744-7658.
PUB. COUNTRY: England; United Kingdom
DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)
LANGUAGE: English
FILE SEGMENT: NONMEDLINE; PUBMED-NOT-MEDLINE
ENTRY MONTH: 200507
ENTRY DATE: Entered STN: 6 Jul 2005
Last Updated on STN: 22 Jul 2005
Entered Medline: 21 Jul 2005

AB Risperidone (Risperdal) is a benzisoxazole derivative with a high affinity for serotonin 5-HT2 and dopamine D2 receptors, and some affinity for alpha- adrenergic, histamine H1 and dopamine D1 receptors. It has no anticholinergic effects. Early studies demonstrated risperidone to be an effective medication for psychotic symptoms, probably more so than the older neuroleptics for both positive and negative symptoms. At clinically effective doses, risperidone causes no more extrapyramidal side-effects (EPS) than placebo; at higher doses EPS frequency increases in a dose-dependent manner. Since it became available in 1994, extensive experience with the drug supports favourable early

impressions of efficacy and tolerability. Minimal sedation, relatively little weight gain and absence of anticholinergic manifestations contribute to the relative tolerability of risperidone as compared to older neuroleptics. However, risperidone is associated with hyperprolactinaemia which can result in amenorrhoea and sexual dysfunction. Compared to older neuroleptics, pharmacoeconomic studies have shown that use of risperidone is associated with reduced hospitalisation and direct cost savings. A recent study found equivalent efficacy between risperidone and clozapine for treatment-resistant patients. Two studies comparing risperidone and olanzapine have yielded positive but conflicting findings. The overall positive experience with risperidone has resulted in the drug being widely recommended as a first line treatment option for psychoses.

L15 ANSWER 2 OF 14 MEDLINE on STN
ACCESSION NUMBER: 2001036398
DOCUMENT NUMBER: Pubmed ID: 1099309
TITLE: Risperidone: a review of its use in the management of the behavioural and psychological symptoms of dementia.
AUTHOR: Bhana N; Spencer C M
CORPORATE SOURCE: Adis International Limited, Mairangi Bay, Auckland, New Zealand.. demall@adis.co.nz
SOURCE: Drugs & aging, (2000 Jun) Vol. 16, No. 6, pp. 451-71. Ref: 74
Journal code: 9102074. ISSN: 1170-229X.

PUB. COUNTRY: New Zealand
DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)
LANGUAGE: English
FILE SEGMENT: Priority Journals
ENTRY MONTH: 200011
ENTRY DATE: Entered STN: 22 Mar 2001
Last Updated on STN: 22 Mar 2001
Entered Medline: 30 Nov 2000

AB Risperidone is a benzisoxazole derivative which has proven efficacy against the positive and negative symptoms of schizophrenia. It has more recently been investigated and shown efficacy as a treatment for the behavioural and psychological symptoms associated with dementia in the elderly. Risperidone has pharmacological properties resembling those of the atypical antipsychotic clozapine and an improved tolerability profile compared with the conventional antipsychotic haloperidol. Risperidone has antagonistic activity primarily at serotonin 5-HT2A and dopamine D2 receptors. In the first 2 large, well controlled trials of an antipsychotic agent used in the treatment of elderly patients with Alzheimer's dementia, vascular dementia or mixed dementia, risperidone 1 mg/day was at least as effective as haloperidol and superior to placebo, as assessed by the rating scales for global behaviour, aggression and psychosis. In extension phases of the 2 trials, clinical benefits were maintained for treatment periods of up to 1 year, with an incidence rate of tardive dyskinesia (2.6%) one-tenth of that seen with conventional antipsychotics. Risperidone, administered at a low dosage of 1 mg/day was associated with fewer extrapyramidal symptoms compared with haloperidol in elderly patients. Risperidone was well tolerated with no clinically relevant abnormalities in laboratory tests, vital signs or electrocardiogram results. Conclusion: The efficacy of risperidone has been demonstrated in the treatment of the behavioural and psychological symptoms associated with dementia in the elderly. Preliminary results from 1-year extension studies confirm the favourable efficacy and tolerability profile of risperidone 1 mg/day. Although head to head studies with other atypical antipsychotic agents are required and the long term use of the drug requires clarification, risperidone represents a generally well tolerated and effective treatment in the management of dementia-associated behavioural and psychological symptoms in the elderly.

L15 ANSWER 3 OF 14 MEDLINE on STN

199187310 MEDLINE
 PUBMED ID: 10087034
 S-16924, a novel, potential antipsychotic with marked serotonin_{1A} agonist properties. IV. A drug discrimination comparison with clozapine.
 AUTHOR: Millan M J; Schreiber R; Monneyron S; Denorme B; Melon C; Queriaux J; Scheyne A
 CORPORATE SOURCE: Institut de Recherches Servier, Centre de Recherches de Croissy, Psychopharmacology Department, Croissy-sur-Seine, Paris, France.
 SOURCE: The Journal of pharmacology and experimental therapeutics, (1999 Apr) Vol. 289, No. 1, pp. 427-36.
 JOURNAL CODE: 0376362. ISSN: 0022-3565.
 PUB. COUNTRY: United States
 DOCUMENT TYPE: (COMPARATIVE STUDY)
 LANGUAGE: English
 ENTRY MONTH: Priority Journals
 ENTRY DATE: 199904
 Entered STN: 4 May 1999
 Last Updated on STN: 18 Jan 2003
 Entered Medline: 21 Apr 1999

AB The novel benzodioxypyrrolidine (S-16924) displays a clozapine-like profile of interaction with multiple monoaminergic receptors, in addition to potent agonist activity at serotonin (5-HT_{1A}) receptors. S-16924 (2.5 mg/kg i.p.) and clozapine (5.0 mg/kg i.p.) generated robust discriminative stimuli (DS) and displayed full mutual generalization. The D4 antagonists L-745,870 and S-18126, the D1/D5 antagonist SCH-39166, and the D3 antagonist S-14297 showed at most partial generalization to S-16924 and clozapine. The D2/D3 antagonist raclopride fully generalized to S-16924, but only partially generalized to clozapine. The 5-HT_{2A} antagonist MDL-100, 907 fully generalized to S-16924 and two further 5-HT_{2A} antagonists, fananserin and SR-46349, showed partial generalization. However, MDL-100, 907, fananserin, and SR-46349 showed less pronounced generalization to clozapine. Similarly, the 5-HT_{2C} antagonists SB-200,646 and SB-206,553 more markedly generalized to S-16924 than to clozapine. The 5-HT_{1A} receptor agonist (7-)-8-dihydroxy-2-(di-n-propylamino) tetralin generalized fully to S-16924 but not to clozapine. Full generalization was obtained to both S-16924 and clozapine for the clozapine congeners, olanzapine and quetiapine. In distinction, the benzisoxazole, risperidone, and the phenylindole, sertindole, weakly generalized to S-16924 and clozapine. However, the benzisoxazole ziprasidone, which possesses 5-HT_{1A} agonist properties, generalized fully to S-16924 but not to clozapine. Finally, the muscarinic antagonist scopolamine generalized fully to clozapine and partially to S-16924. In conclusion, S-16924 and clozapine display both communalities and differences in their "compound" DS; this likely reflects their respective complex patterns of interaction with multiple monoaminergic receptors. Although no specific receptor was identified as underlying the clozapine DS, 5-HT_{1A} agonist as well as D2 and 5-HT_{2A/2C} antagonist properties contribute to the S-16924 DS.

L15 ANSWER 4 OF 14 MEDLINE on STN
 ACCESSION NUMBER: 97308342
 DOCUMENT NUMBER: MEDLINE
 PUBMED ID: 9165568
 TITLE: Neuroleptic malignant syndrome with risperidone.
 AUTHOR: Gleason P P; Conigliaro R L
 CORPORATE SOURCE: Department of Pharmaceutical Sciences, School of Pharmacy, University of Pittsburgh, Pennsylvania, USA.
 SOURCE: Pharmacotherapy, (1997 May-Jun) Vol. 17, No. 3, pp. 617-21.
 JOURNAL CODE: 8111305. ISSN: 0277-0008.
 PUB. COUNTRY: United States
 DOCUMENT TYPE: (CASE REPORTS)
 LANGUAGE: English
 ENTRY MONTH: Journal; Article; (JOURNAL ARTICLE)

FILE SEGMENT: Priority Journals
 ENTRY MONTH: 199707
 ENTRY DATE: Entered STN: 24 Jul 1997
 Last Updated on STN: 24 Jul 1997
 Entered Medline: 15 Jul 1997

AB Neuroleptic malignant syndrome is thought to be a result of dopamine D2 receptor blockade in the striatum of the basal ganglia. Risperidone, a benzisoxazole derivative antipsychotic, has high serotonin 5-HT₂ receptor blockade and dose-related D2 receptor blockade. The high ratio is believed to impart the low frequency of extrapyramidal symptoms with risperidone at low dosages. With this low frequency of extrapyramidal symptoms, it was thought the frequency of neuroleptic malignant syndrome might also be lowered. A 73-year-old woman developed neuroleptic malignant syndrome after monotherapy with risperidone. The syndrome reversed after discontinuing risperidone and starting treatment with dantrolene and bromocriptine. It appears that the protection from extrapyramidal side effects observed with risperidone does not ensure protection from neuroleptic malignant syndrome.

L15 ANSWER 5 OF 14 MEDLINE on STN
 ACCESSION NUMBER: 96142226
 DOCUMENT NUMBER: MEDLINE
 PUBMED ID: 8543544
 TITLE: Risperidone as a treatment for Tourette's syndrome.
 AUTHOR: Bruun R D; Budman C L
 CORPORATE SOURCE: Cornell University Medical School, New York, N.Y., USA.
 SOURCE: The Journal of clinical psychiatry, (1996 Jan) Vol. 57, No. 1, pp. 29-31.
 JOURNAL CODE: 7801243. ISSN: 0160-6689.
 PUB. COUNTRY: United States
 DOCUMENT TYPE: (CLINICAL TRIAL)
 LANGUAGE: English
 ENTRY MONTH: Priority Journals
 ENTRY DATE: 199602
 Entered STN: 27 Feb 1996
 Last Updated on STN: 27 Feb 1996
 Entered Medline: 9 Feb 1996

AB BACKGROUND: An open-label trial was performed to assess the efficacy and safety of risperidone, a benzisoxazole derivative with potent D2 and 5-HT₂ antagonism, for treatment of Tourette's syndrome. METHOD: Thirty-eight patients with Tourette's syndrome volunteered to take risperidone for treatment of their tics. All patients had failed to respond adequately to conventional treatments (with neuroleptics such as haloperidol and/or with the alpha 2-adrenergic agonist clonidine) or had suffered from intolerable side effects from such treatments. Patients were rated for tic severity by the Yale Global Tic Severity Scale (YGTSS) before treatment and after 1 month of treatment with risperidone. Patients were monitored carefully for side effects and clinical response. RESULTS: Of the 38 patients, 8 discontinued risperidone treatment before the end of the trial because of intolerable side effects. At the end of the 4-week trial, 22 patients (58%) were improved, 7 patients (18%) had no appreciable change in their symptoms, and 1 patient (3%) had a documented worsening of tics. Doses of risperidone at the end of the trial ranged from 0.5 mg to 9 mg/day (mean = 2.7 mg/day). CONCLUSION: This open clinical trial suggests that risperidone may be a promising alternative to conventional medications used for treating the symptoms of Tourette's syndrome.

L15 ANSWER 6 OF 14 MEDLINE on STN
 ACCESSION NUMBER: 95348446
 DOCUMENT NUMBER: MEDLINE
 PUBMED ID: 7542676
 TITLE: A pharmacological, pharmacokinetic and clinical overview of risperidone, a new antipsychotic that blocks serotonin 5-HT₂ and dopamine D2 receptors.
 AUTHOR: He H; Richardson J S

CORPORATE SOURCE: College of Pharmacy, College of Medicine, University of Saskatchewan, Saskatoon, Canada.
SOURCE: International clinical psychopharmacology, (1995 Mar) Vol. 10, No. 1, pp. 19-30. Ref: 78
 Journal code: 8609061. ISSN: 0268-1315.
PUB. COUNTRY: ENGLAND: United Kingdom
DOCUMENT TYPE: Journal Article; (JOURNAL ARTICLE)
 General Review; (REVIEW)
LANGUAGE: English
FILE SEGMENT: Priority Journals
ENTRY MONTH: 199508
ENTRY DATE: Entered STN: 11 Sep 1995
 Last Updated on STN: 29 Jan 1996
 Entered Medline: 30 Aug 1995

AB Risperidone is a benzisoxazole derivative with antipsychotic activity that is chemically unrelated to other currently available antipsychotic agents. Its neuropharmacological properties, characterized by potent central antagonism of both serotonin 5-HT₂ and dopamine D₂ receptors, also differ from those of most other antipsychotic drugs. The pharmacokinetics of risperidone are well understood, having been studied in healthy subjects as well as in psychotic patients. The absolute oral bioavailability of risperidone is nearly 70%, and after oral administration, it is rapidly absorbed with the plasma level reaching a peak at about 1 h. 9-Hydroxyrisperidone, one of the metabolites of risperidone, is equally active with the parent compound and so the clinical activity of a dose of risperidone is due to the combined actions of both moieties. The plasma concentrations of risperidone and its active metabolite remain dose proportional even at doses exceeding the therapeutic range. In clinical trials with chronic schizophrenia patients, risperidone has an overall therapeutic activity comparable with that of haloperidol, but at doses that produce similar improvements in the positive symptoms of schizophrenia, risperidone has a greater effect on the negative symptoms and produces less extrapyramidal side effects than does haloperidol. However, additional controlled clinical studies are needed before the claims that risperidone is therapeutically superior to haloperidol can be considered to be established firmly. Although risperidone is effective in acute schizophrenia and in non-treatment-resistant schizophrenics, studies adequately comparing risperidone with clozapine in treatment-resistant schizophrenic patients remain to be published. In addition, risperidone has been reported to be of value in patients with schizodepressive disorders. The clinical success of risperidone suggests that the development of compounds with selective affinity for 5-HT₂ or other serotonin receptors may result in even further improvements in the pharmacotherapy of psychiatric disorders.

L15 ANSWER 7 OF 14 MEDLINE on STN
ACCESSION NUMBER: 95222523 MEDLINE
DOCUMENT NUMBER: PubMed ID: 7707315
TITLE: 3-[[[Aryloxy]alkyl]piperidinyl]-1,2-benzisoxazoles as D₂/5-HT₂ antagonists with potential atypical antipsychotic activity: antipsychotic profile of alogaperidone (HP 873).
AUTHOR: Strupczewski J T; Bordeaux K J; Chiang Y; Glankowski E J; Conway P G; Corbett R; Hartman H B; Szelezak M R; Wilmot C A; Helsley G C
CORPORATE SOURCE: Chemical Research Department, Hoechst-Roussel Pharmaceuticals Inc, Somerville, New Jersey 08876, USA.
SOURCE: Journal of medicinal chemistry, (1995 Mar 31) Vol. 38, No. 7, pp. 1119-31.
 Journal code: 9716531. ISSN: 0022-2623.
PUB. COUNTRY: United States
DOCUMENT TYPE: (IN VITRO)
LANGUAGE: Journal Article; (JOURNAL ARTICLE)
FILE SEGMENT: English
 Priority Journals

ENTRY MONTH: 199505
ENTRY DATE: Entered STN: 18 May 1995
 Last Updated on STN: 18 May 1995
 Entered Medline: 11 May 1995

AB A series of 3-[[[aryloxy]alkyl]piperidinyl]-1,2-benzisoxazoles was synthesized and evaluated as potential antipsychotic D₂/5-HT₂ antagonists. Most of these compounds showed potent antipsychotic-like activity in an apomorphine-induced climbing mouse paradigm, with many also showing preferential mesolimbic activity, as indicated by their weaker effects in an apomorphine-induced stereotypy model. In receptor binding assays, many displayed a moderate affinity for the D₂ receptor coupled with a significantly greater affinity for the 5-HT₂ receptor: a property that has been suggested as necessary for atypicality. From this series, compound 45, 1-[[4-[3-[4-(6-fluoro-1,2-benzisoxazol-3-yl)-1-piperidinyl]propoxy]-3-methoxyphenyl]ethanone (iloperidone, HP 873), was further evaluated in a battery of in vivo and in vitro assays. This compound showed a 300-fold greater potency in inhibition of climbing than in inhibition of stereotypy or induction of catalepsy, and when evaluated chronically in an electrophysiological model, 45 caused a depolarization blockade of dopamine neurons in the A10 area of the rat brain but not in the A9 area. Additionally, it showed positive activity in a social interaction paradigm, suggesting potential efficacy against sociality, a component of the negative symptoms of schizophrenia. In chronic ex vivo studies, 45, similar to clozapine, caused a down regulation of 5-HT₂ receptors but had no effect on the number of D₂ receptors. Compound 45 is currently undergoing clinical evaluation.

L15 ANSWER 8 OF 14 MEDLINE on STN
ACCESSION NUMBER: 95148787 MEDLINE
DOCUMENT NUMBER: PubMed ID: 7531352
TITLE: Regional brain distribution of risperidone and its active metabolite 9-hydroxy-risperidone in the rat.
AUTHOR: van Beijsterveldt L E; Geerts R J; Leyssen J E; Megens A A; Van den Eynde H M; Meuldermans W E; Heykants J J
CORPORATE SOURCE: Department of Drug Metabolism and Pharmacokinetics, Janssen Research Foundation, Beerse, Belgium.
SOURCE: Psychopharmacology, (1994 Feb) Vol. 114, No. 1, pp. 53-62.
 Journal code: 7608025. ISSN: 0033-3158.
PUB. COUNTRY: GERMANY: Germany, Federal Republic of
DOCUMENT TYPE: Journal Article; (JOURNAL ARTICLE)
LANGUAGE: English
FILE SEGMENT: Priority Journals
ENTRY MONTH: 199503
ENTRY DATE: Entered STN: 16 Mar 1995
 Last Updated on STN: 29 Jan 1996
 Entered Medline: 8 Mar 1995

AB Risperidone is a new benzisoxazole antipsychotic. 9-Hydroxy-risperidone is the major plasma metabolite of risperidone. The pharmacological properties of 9-hydroxy-risperidone were studied and appeared to be comparable to those of risperidone itself, both in respect of the profile of interactions with various neurotransmitters and its potency, activity, and onset and duration of action. The absorption, plasma levels and regional brain distribution of risperidone, metabolically formed 9-hydroxy-risperidone and total radioactivity were studied in the male Wistar rat after single subcutaneous administration of radiolabelled risperidone at 0.02 mg/kg. Concentrations were determined by HPLC separation, and off-line determination of the radioactivity with liquid scintillation counting. Risperidone was well absorbed. Maximum plasma concentrations were reached at 0.5-1 h after subcutaneous administration. Plasma concentrations of 9-hydroxy-risperidone were higher than those of risperidone from 2h after dosing. In plasma, the apparent elimination half-life of risperidone was 1.0 h, and mean residence times were 1.5 h for risperidone and 2.5 h for its 9-hydroxy metabolite. Plasma levels of the radioactivity increased dose

proportionally between 0.02 and 1.3 mg/kg. Risperidone was rapidly distributed to brain tissues. The elimination of the radioactivity from the frontal cortex and striatum-brain regions with high concentrations of 5-HT₂ or dopamine-D₂ receptors--became more gradual with decreasing dose levels. After a subcutaneous dose of 0.02 mg/kg, the ED50 for central 5-HT₂ antagonism in male rats, half-lives in frontal cortex and striatum were 3-4 h for risperidone, whereas mean residence times were 4-6 h for risperidone and about 12 h for 9-hydroxy-risperidone. These half-lives and mean residence times were 3-5 times longer than in plasma and in cerebellum, a region with very low concentrations of 5-HT₂ and D₂ receptors. Frontal cortex and striatum to plasma concentration ratios increased during the experiment. The distribution of 9-hydroxy-risperidone to the different brain regions, including frontal cortex and striatum, was more limited than that of risperidone itself. This indicated that 9-hydroxy-risperidone contributes to the in vivo activity of risperidone, but to a smaller extent than would be predicted from plasma levels. AUCs of both active compounds in frontal cortex and striatum were 10-18 times higher than those in cerebellum. No retention of metabolites other than 9-hydroxy-risperidone was observed in any of the brain regions investigated.

L15 ANSWER 9 OF 14 MEDLINE on STN
 ACCESSION NUMBER: 95023318 MEDLINE
 DOCUMENT NUMBER: PubMed ID: 7524043
 TITLE: Risperidone.
 AUTHOR: Cohen L J
 CORPORATE SOURCE: College of Pharmacy, University of Oklahoma, Oklahoma City 73190.
 SOURCE: Journal code: 8111305. ISSN: 0277-0008.
 Ref: 62
 United States
 (COMPARATIVE STUDY)
 Journal; Article; (JOURNAL ARTICLE)
 English
 Priority Journals
 199411
 Entered STN: 22 Dec 1994
 Last Updated on STN: 29 Jan 1996
 Entered Medline: 21 Nov 1994

AB Risperidone, a benzisoxazole derivative, is a novel antipsychotic agent that has an extremely strong binding affinity for serotonin 5-HT₂ receptors, a strong binding affinity for dopamine D₂ receptors, and a high affinity for alpha 1- and alpha 2-adrenergic receptors and histamine H₁ receptors. Its affinity for serotonin receptors is approximately 200 times greater than that of haloperidol, and its dopamine antagonistic potency is comparable to that of haloperidol. Its major metabolite, 9-hydroxyrisperidone, has similar pharmacologic activity, and thus the parent compound and metabolite form the active antipsychotic moiety. Clinical trials demonstrate that risperidone is an effective antipsychotic agent that improves negative as well as positive symptoms of schizophrenia. At recommended dosages, the frequency of extrapyramidal side effects is no greater than that seen with placebo. The drug appears to be an advance in the treatment of psychoses.

L15 ANSWER 10 OF 14 MEDLINE on STN
 ACCESSION NUMBER: 94334885 MEDLINE
 DOCUMENT NUMBER: PubMed ID: 7914536
 TITLE: Benzisoxazole- and benzisothiazole-3-carboxamides as potential atypical antipsychotic agents.
 AUTHOR: Hrib N J; Jurcak J G; Burgher K L; Conway P G; Hartman H B; Kernan L J; Roehr J E; Woods A T
 CORPORATE SOURCE: Neuroscience Strategic Business Unit, Hoechst-Roussel Pharmaceuticals, Inc., Somerville, New Jersey 08876.

SOURCE: Journal of medicinal chemistry, (1994 Jul 22) Vol. 37, No. 15, pp. 2308-14.
 Journal code: 9716531. ISSN: 0022-2623.

PUB. COUNTRY: United States
 DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)
 LANGUAGE: English
 FILE SEGMENT: Priority Journals
 ENTRY MONTH: 199409
 ENTRY DATE: Entered STN: 20 Sep 1994
 Last Updated on STN: 6 Feb 1995
 Entered Medline: 12 Sep 1994

AB A series of benzisoxazole- and benzisothiazole-3-carboxamides has been prepared and tested for potential antipsychotic activity. In general, the compounds showed an affinity for dopamine D₂ and serotonin 5HT_{2A} and 5HT_{1A} receptors. Several members of this series have demonstrated activity in animal models predictive of potential antipsychotic activity. In addition, compounds 18, 19, 22, 27, 28, 43, and 44 have also shown a potential for reduced EPS liability as suggested by the ratio of activity seen in mesolimbic-mediated vs nigrostriatal-mediated behavioral assays.

L15 ANSWER 11 OF 14 MEDLINE on STN
 ACCESSION NUMBER: 94046936 MEDLINE
 DOCUMENT NUMBER: PubMed ID: 7901415
 TITLE: Examination of the D₂/5-HT₂ affinity ratios of resolved 5,6,7,8,9,10-hexahydro-7,10-iminocyclohept[b]indoles: an enantioselective approach toward the design of potential atypical antipsychotics.
 AUTHOR: Mewshaw R E; Abreu M E; Silverman L S; Mathew R M; Tiffany C W; Bailey M A; Karbon E W; Feikany J W; Kaiser C
 CORPORATE SOURCE: Department of Medicinal Chemistry, Scios Nova Inc., Baltimore, Maryland 21224-6522.
 SOURCE: Journal of medicinal chemistry, (1993 Oct 15) Vol. 36, No. 21, pp. 3073-6.
 Journal code: 9716531. ISSN: 0022-2623.

PUB. COUNTRY: United States
 DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)
 LANGUAGE: English
 FILE SEGMENT: Priority Journals
 ENTRY MONTH: 199311
 ENTRY DATE: Entered STN: 17 Jan 1994
 Last Updated on STN: 3 Feb 1997
 Entered Medline: 26 Nov 1993

AB Enantiomers of several N-substituted 5,6,7,8,9,10-hexahydro-7,10-iminocyclohept[b]indoles were obtained by the resolution of 5,6,7,8,9,10-hexahydro-7,10-iminocyclohept[b]indole and N-alkylation. These, as well as the racemates, were evaluated for their affinity for the 5-HT₂ and D₂ receptors. Those compounds possessing the 7S,10R stereochemistry were consistently recognized by the 5-HT₂ and D₂ receptors as the eutomer. 2-Fluoro-11-[4-(4-fluorophenyl)-4-oxobutyl]-5,6,7,8,9,10-hexahydro-7S,10 R-iminocyclohept[b]indole [(7S,10R)-8] had the highest affinity for the 5-HT₂ receptor (K_i = 0.80 nM), while its diastomer (7R,10S)-8 was the most selective member of this class of bridged gamma-carbolines (D₂ 7S-HT₂ = 562). Incorporation of a benzoyl or isosteric benzisoxazole moiety tethered by a four-carbon spacer to a bridged gamma-carboline nucleus, possessing the 7S,10R absolute configuration, produced high affinity ligands for the 5-HT₂ and D₂ receptors.

L15 ANSWER 12 OF 14 MEDLINE on STN
 ACCESSION NUMBER: 93267591 MEDLINE
 DOCUMENT NUMBER: PubMed ID: 8496917
 TITLE: Bridged gamma-carbolines and derivatives possessing selective and combined affinity for 5-HT₂ and D₂

receptors.
 Mewshaw R E; Silverman L S; Mathew R M; Kaiser C; Sherrill R G; Cheng M; Tiffany C W; Karbon E W; Bailey M A; Borosky S A; +
 Scios Nova Inc., Baltimore, Maryland 21224-6522.
 Journal of medicinal chemistry, (1993 May 14) Vol. 36, No. 10, pp. 1488-95.
 Journal code: 9716531. ISSN: 0022-2623.
 United States
 Journal; Article; (JOURNAL ARTICLE)
 English
 Priority Journals
 199306
 Entered STN: 2 Jul 1993
 Last Updated on STN: 2 Jul 1993
 Entered Medline: 22 Jun 1993
 AB A series of 5,6,7,8,9,10-hexahydro-7,10-iminocyclohept(b)indoles and 6,7,8,9,10,11-hexahydro-7,11-imino-5H-cyclooct(b)indoles was prepared. Structural modifications of the lead compound, 11-[4-(4-fluorobenzoyl)propyl]-5,6,7,8,9,10-hexahydro-7,10-iminocyclohept(b)indole (5, Ki = 0.82 nM vs [3H]ketanserin) enabled the identification of the functionality necessary for high affinity at serotonin 5-HT2 and dopamine D2 receptors in ligand binding studies. The indole ring, as well as the benzoyl or isosteric benzisoxazole moiety, were essential for high affinity. Variations of the length of the side chains resulted in ligands having either selective affinity for the 5-HT2 receptor or a combination of 5-HT2 and D2 affinity. In vivo binding studies were performed on selected members in this series. The most potent member, 2-fluoro-11-[4-(4-fluorobenzoyl)butyl]-5,6,7,8,9,10-hexahydro-7,10-iminocyclohept(b)indole (36) had an ED50 of < 1 mg/kg at the 5-HT2 and D2 receptors following oral administration.

L15 ANSWER 13 OF 14 MEDLINE on STN
 ACCESSION NUMBER: 91157659
 DOCUMENT NUMBER: 2127339
 TITLE: [Desarrollo de nuevas drogas antipsicoticas. El desarrollo de nuevas drogas antipsicoticas. Vanden Bussche G; Gelders Y G; Heylen S L Clinical Research and Development Department, Janssen Research Foundation, Beerse, Belgium. Acta psiquiatrica y psicologica de America latina. (1990 Jan-Jun) Vol. 36, No. 1-2, pp. 13-25. Journal code: 0373060. ISSN: 0001-6896. Argentina (ENGLISH ABSTRACT) Journal; Article; (JOURNAL ARTICLE) Spanish Priority Journals 199104
 Entered STN: 28 Apr 1991
 Last Updated on STN: 28 Apr 1991
 Entered Medline: 8 Apr 1991
 AB As far as schizophrenia is concerned, therapeutic effects of neuroleptics based on brain-located dopamine receptor blockers are taken for granted. It is also admitted, however, that classical neuroleptics have inconveniences, namely: Their relative lack of effect on negative symptoms, and their liability to induce extrapyramidal symptoms (EPS). Pimaperone-based clinical studies evidenced that an antagonist combining serotonin 5-HT2, and dopamine D2 was successful in the treatment of schizophrenia--which could be clearly observed in (a) anti-austic effects, (b) regulating disrupted sleep-wake rhythms, and (c) a lesser tendency to EPS. Seropone-based studies--a compound with a comparable pharmacological profile--confirmed the above observations. Until, however, the synthesis of ritanerlin--a specific, and selective antagonistic receptor--was not achieved, no exact implication of 5-HT2

AUTHOR:
 CORPORATE SOURCE:
 SOURCE:
 PUB. COUNTRY:
 DOCUMENT TYPE:
 LANGUAGE:
 FILE SEGMENT:
 ENTRY MONTH:
 ENTRY DATE:

antagonist in psychopharmacological treatments of schizophrenia could be explored further. Indeed, double-blind trials evidenced a remarkable improvement in negative as well as extrapyramidal symptoms. Since a monotherapy appeared as undeniably called for in the treatment of schizophrenia, the next logical step to be taken was selecting a compound with a central antagonism comparable to ritanerlin's, and a central D2 antagonism comparable to haloperidol's. Among a chemical range of benzisoxazole derivatives, risperidone was thus selected. The first double-blind trials on chronic schizophrenic patients seem indeed to confirm that this substance is likely to get over the above mentioned inconveniences, so typical of classical neuroleptics.

L15 ANSWER 14 OF 14 MEDLINE on STN
 ACCESSION NUMBER: 88153393
 DOCUMENT NUMBER: 2450200
 TITLE: Pharmacology of risperidone (R 64 766), a new antipsychotic with serotonin-S2 and dopamine-D2 antagonistic properties.
 AUTHOR: Janssen P A; Niemegeers C J; Awouters F; Schellekens K H; Megens A A; Meert T F
 CORPORATE SOURCE: Department of Pharmacology, Janssen Research Foundation, Beerse, Belgium.
 SOURCE: The Journal of pharmacology and experimental therapeutics, (1988 Feb) Vol. 244, No. 2, pp. 685-93. Journal code: 0376362. ISSN: 0022-3565. United States
 Journal; Article; (JOURNAL ARTICLE)
 English
 Priority Journals
 198804
 Entered STN: 8 Mar 1990
 Last Updated on STN: 6 Feb 1995
 Entered Medline: 12 Apr 1988
 AB Comparative studies of the benzisoxazole derivative risperidone (R 64 766) were made with ritanerlin, a selective centrally acting serotonin-S2 antagonist and with haloperidol, a selective centrally acting dopamine-D2 antagonist. Risperidone like ritanerlin shows activity in all tests related to serotonin-S2 antagonism, but at even lower doses (peripheral S2-antagonism at 0.001 mg/kg, central S2-antagonism at 0.014 mg/kg). Like haloperidol, risperidone shows activity in all tests related to dopamine-D2 antagonism; central nervous system controlled functions, including the induction of catalepsy, are relatively much less affected by risperidone. Qualitatively, risperidone is a mixed serotonin-dopamine antagonist. Quantitatively, its study in dogs reveals potent dopamine-D2 antagonistic activity with excellent p.o. bioavailability and a relatively long duration of action. From the obtained pharmacological data, risperidone could be expected to possess the complementary clinical effects of a ritanerlin-like serotonin-S2 and an haloperidol-like dopamine-D2 antagonist. Serotonin-S2 antagonism may improve the quality of sleep, reduce negative and affective symptoms in schizophrenic patients and decrease extrapyramidal symptoms induced by classical neuroleptics. Because risperidone is a dopamine-D2 antagonist, antidelusional, antihallucinatory and antinamic actions are expected. The first clinical studies indicate that two additional therapeutic targets, which are not reached with classical neuroleptics, may be obtained with risperidone in the monotherapy of schizophrenia and related disorders: very important contact and mood-elevating properties and extrapyramidal symptoms-free maintenance therapy.

==>

----Logging off of STN----

NEWS 30 APR 02 JICST-EPLUS removed from database clusters and STN

NEWS EXPRESS NOVEMBER 10 CURRENT WINDOWS VERSION IS V8.01c. CURRENT
MACINTOSH VERSION IS V6.0c(ENG) AND V6.0Jc(JP),
AND CURRENT DISCOVER FILE IS DATED 25 SEPTEMBER 2006.

NEWS HOURS STN Operating Hours Plus Help Desk Availability
NEWS LOGIN Welcome Banner and News Items
NEWS IPC8 For general information regarding STN implementation of IPC 8
NEWS X25 X.25 communication option no longer available

Enter NEWS followed by the item number or name to see news on that
specific topic.

All use of STN is subject to the provisions of the STN Customer
agreement. Please note that this agreement limits use to scientific
research. Use for software development or design or implementation
of commercial gateways or other similar uses is prohibited and may
result in loss of user privileges and other penalties.

***** STN Columbus *****

FILE 'HOME' ENTERED AT 13:55:13 ON 09 APR 2007

=> file medline

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
	0.21	0.21

FULL ESTIMATED COST

FILE 'MEDLINE' ENTERED AT 13:55:53 ON 09 APR 2007

FILE LAST UPDATED: 7 Apr 2007 (20070407/UP). FILE COVERS 1950 TO DATE.

This file contains CAS Registry Numbers for easy and accurate
substance identification.

=> s 5ht1b or 5-ht1b or 5ht-1bh
137 5HT1B
2352638 5

1858 HT1B
1855 5-HT1B
(5(W)HT1B)

2604 5HT
2 1BH
0 5HT-1BH
(5HT(W)1BH)

L1 1953 5HT1B OR 5-HT1B OR 5HT-1BH

=> s 5ht1b or 5-ht1b or 5ht-1b
137 5HT1B
2352638 5

1858 HT1B
1855 5-HT1B
(5(W)HT1B)

2604 5HT
13798 1B
55 5HT-1B
(5HT(W)1B)

L2 1984 5HT1B OR 5-HT1B OR 5HT-1B

=> s l2 and antagonist?

L3 575230 ANTAGON?
1099 L2 AND ANTAGON?

=> s l3 and new

=>
Executing the logoff script...

=> LOG Y

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
	16.41	47.87

FULL ESTIMATED COST

STN INTERNATIONAL LOGOFF AT 13:24:47 ON 09 APR 2007

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:sssptal623zct

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

***** Welcome to STN International *****

NEWS 1 Web Page URLs for STN Seminar Schedule - N. America

NEWS 2 "Ask CAS" for self-help around the clock

NEWS 3 DEC 18 CA/CAPLUS pre-1967 chemical substance index entries enhanced

NEWS 4 DEC 18 with preparation role

NEWS 5 DEC 18 CA/CAPLUS patent kind codes updated

NEWS 6 DEC 18 MARPAT to CA/CAPLUS accession number crossover limit increased to 50,000

NEWS 7 DEC 18 MEDLINE updated in preparation for 2007 reload

NEWS 8 JAN 08 CA/CAPLUS enhanced with more pre-1907 records

NEWS 9 JAN 16 CHEMLIST enhanced with New Zealand Inventory of Chemicals

NEWS 10 JAN 16 CA/CAPLUS Company Name Thesaurus enhanced and reloaded

NEWS 11 JAN 16 IPC version 2007.01 thesaurus available on STN

NEWS 12 JAN 16 WPIDS/WPIINDEX/WPIX enhanced with IPC 8 reclassification data

NEWS 13 JAN 22 CA/CAPLUS updated with revised CAS roles

NEWS 14 JAN 22 PHAR reloaded with new search and display fields

NEWS 15 JAN 29 CAS Registry Number crossover limit increased to 300,000 in multiple databases

NEWS 16 FEB 15 PATDPASPC enhanced with Drug Approval numbers

NEWS 17 FEB 15 RUSSIAPAT enhanced with pre-1994 records

NEWS 18 FEB 23 KOREAPAT enhanced with IPC 8 features and functionality

NEWS 19 FEB 26 MEDLINE reloaded with enhancements

NEWS 20 FEB 26 EMBASE enhanced with Clinical Trial Number field

NEWS 21 FEB 26 TOXCENTER enhanced with reloaded MEDLINE

NEWS 22 FEB 26 IFICDB/IFIPAT/IFIUDB reloaded with enhancements

NEWS 23 FEB 26 CAS Registry Number crossover limit increased from 10,000 to 300,000 in multiple databases

NEWS 24 MAR 15 WPIDS/WPIX enhanced with new FRAGHITSTR display format

NEWS 25 MAR 16 CASREACT coverage extended

NEWS 26 MAR 20 MARPAT now updated daily

NEWS 27 MAR 22 LNPFI reloaded

NEWS 28 MAR 30 RDISCLOSURE reloaded with enhancements

NEWS 29 MAR 30 INPADOCDB will replace INPADOC on STN

1000180 NEW
5806 NEWS
1005218 NEW
(NEW OR NEWS)
L4 67 L3 AND NEW
=> s 14 and 2003/Py
569314 2003/Py
(20030000-20039999/Py)
L5 4 L4 AND 2003/Py
=> d 1-4 ibib abs

L5 ANSWER 1 OF 4 MEDLINE on STN
ACCESSION NUMBER: 2004418030 MEDLINE
DOCUMENT NUMBER: PubMed ID: 15320857
TITLE: Migraine: pathophysiology, pharmacology, treatment and future trends.
AUTHOR: Villalon Carlos M; Centurion David; Valdivia Luis Felipe; de Vries Peter; Saxena Pramod R
CORPORATE SOURCE: Departamento de Farmacobiología, CINVESTAV-IPN, Czada. de los Tenorios 235, Col. Granjas Coapa, Deleg. Tlalpan, CP 14300, Mexico DF, Mexico. carlos.villalon@infotel.net.mx
SOURCE: Current vascular pharmacology, (2003 Mar) Vol. 1, No. 1, pp. 71-84. Ref: 110
Journal code: 101157208. ISSN: 1570-1611.
United Arab Emirates
PUB. COUNTRY: Historical
DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)
(RESEARCH SUPPORT, NON-U.S. GOV'T)
General Review; (REVIEW)
LANGUAGE: English
FILE SEGMENT: Priority Journals
ENTRY MONTH: 200412
ENTRY DATE: Entered STN: 25 Aug 2004
Last Updated on STN: 19 Dec 2004
Entered Medline: 1 Dec 2004

AB Migraine treatment has evolved into the scientific arena, but it seems still controversial whether migraine is primarily a vascular or a neurological dysfunction. Irrespective of this controversy, the levels of serotonin (5-hydroxytryptamine; 5-HT), a vasoconstrictor and a central neurotransmitter, seem to decrease during migraine (with associated carotid vasodilatation) whereas an i.v. infusion of 5-HT can abort migraine. In fact, 5-HT as well as ergotamine, dihydroergotamine and other antimigraine agents invariably produce vasoconstriction in the external carotid circulation. The last decade has witnessed the advent of sumatriptan and second generation triptans (e.g. zolmitriptan, rizatriptan, naratriptan), which belong to a new class of drugs, the 5-HT1B/1D/1F receptor agonists. Compared to sumatriptan, the second-generation triptans have a higher oral bioavailability and longer plasma half-life. In line with the vascular and neurogenic theories of migraine, all triptans produce selective carotid vasoconstriction (via 5-HT1B receptors) and presynaptic inhibition of the trigeminovascular inflammatory responses implicated in migraine (via 5-HT1D/5-HT1F receptors). Moreover, selective agonists at 5-HT1D/5-HT1F (PNU-142633) and 5-HT1F (LY344864) receptors inhibit the trigeminovascular system without producing vasoconstriction. Nevertheless, PNU-142633 proved to be ineffective in the acute treatment of migraine, whilst LY344864 did show some efficacy when used in doses which interact with 5-HT1B receptors. Finally, although the triptans are effective antimigraine agents producing selective cranial vasoconstriction, efforts are being made to develop other effective antimigraine alternatives acting via the direct blockade of vasodilator mechanisms (e.g. antagonists at CGRP receptors, antagonists at 5-HT7 receptors, inhibitors of nitric oxide

biosynthesis, etc). These alternatives will hopefully lead to fewer side effects.

L5 ANSWER 2 OF 4 MEDLINE on STN
ACCESSION NUMBER: 2003200109 MEDLINE
DOCUMENT NUMBER: PubMed ID: 12644890
TITLE: Role of extracellular serotonin levels in the effect of 5-HT1B receptor blockade.
AUTHOR: de Groote Lotte; Klompmaekers Andre A; Olivier Berend; Westenbery Herman G M
CORPORATE SOURCE: Department of Psychiatry, Rudolf Magnus Institute of Neuroscience, University Medical Center Utrecht, Heidelberglaan 100, 3584 CX Utrecht, The Netherlands. pp. 153-8. Electronic Publication: 2003-03-18.
Journal code: 7608025. ISSN: 0033-3158.
Germany; Germany, Federal Republic of
PUB. COUNTRY: Journal; Article; (JOURNAL ARTICLE)
DOCUMENT TYPE: Priority Journals
LANGUAGE: English
FILE SEGMENT: 200310
ENTRY MONTH: Entered STN: 30 Apr 2003
ENTRY DATE: Last Updated on STN: 15 Oct 2003
Entered Medline: 14 Oct 2003

AB The release of serotonin (5-HT) at serotonergic nerve terminals is regulated by 5-HT(1B) autoreceptors. Several studies have reported that the effects of selective 5-HT(1B) receptor inhibitors (SSRIs) on extracellular 5-HT are augmented by 5-HT(1B) receptor antagonists, whereas administration of these antagonists alone do not enhance 5-HT levels. It has been suggested that 5-HT(1B) receptors have low basal endogenous activity and therefore elevated endogenous 5-HT levels are needed to elicit an effect of 5-HT(1B) receptor antagonists. To test this hypothesis, different strategies were used to enhance 5-HT levels in the rat frontal cortex to assess the effects of locally applied NAS-181, a new selective 5-HT(1B) receptor antagonist. Blockade of 5-HT(1B) receptors with NAS-181 dose dependently augmented 5-HT levels when 5-HT levels were enhanced by a SSRI. No additional effect of NAS-181 on 5-HT output was found when 5-HT levels were enhanced by KCl depolarization-induced release or by preventing degradation of 5-HT with the monoamine oxidase inhibitor pargyline. In the presence of fluvoxamine, the increased 5-HT release evoked by KCl depolarization was augmented by NAS-181, supporting the idea that blockade of 5-HT transporters is necessary to measure an effect of 5-HT(1B) receptor blockade. In conclusion, the results provide circumstantial evidence that the effect of a 5-HT(1B) receptor antagonist depends on extracellular 5-HT levels, but strongly suggest that additional 5-HT reuptake inhibition is required to detect any effect of 5-HT(1B) receptor antagonist on 5-HT levels by in vivo microdialysis.

L5 ANSWER 3 OF 4 MEDLINE on STN
ACCESSION NUMBER: 2003132987 MEDLINE
DOCUMENT NUMBER: PubMed ID: 12595948
TITLE: An evaluation of the effect of NAS-181, a new selective 5-HT(1B) receptor antagonist, on extracellular 5-HT levels in rat frontal cortex.
AUTHOR: de Groote Lotte; Klompmaekers Andre A; Olivier Berend; Westenbery Herman G M
CORPORATE SOURCE: Department of Psychiatry, Rudolf Magnus Institute of Neuroscience, University Medical Center Utrecht, Heidelberglaan 100, 3584 CX, Utrecht, The Netherlands. Naunyn-Schmiedeberg's archives of pharmacology, (2003 Feb) Vol. 367, No. 2, pp. 89-94. Electronic Publication: 2003-01-24.
Journal code: 0326264. ISSN: 0028-1298.
Germany; Germany, Federal Republic of
PUB. COUNTRY: Journal; Article; (JOURNAL ARTICLE)
DOCUMENT TYPE: Priority Journals
LANGUAGE: English
FILE SEGMENT: 200310
ENTRY MONTH: Entered STN: 30 Apr 2003
ENTRY DATE: Last Updated on STN: 15 Oct 2003
Entered Medline: 14 Oct 2003

DOCUMENT TYPE: (COMPARATIVE STUDY)
 Journal: Article; (JOURNAL ARTICLE)
 LANGUAGE: English
 FILE SEGMENT: Priority Journals
 ENTRY MONTH: 200312
 ENTRY DATE: Entered STN: 22 Mar 2003
 Last Updated on STN: 24 Dec 2003
 Entered Medline: 23 Dec 2003

AB In the mammalian brain 5-HT_{1B} receptors are present as autoreceptors regulating the release of serotonin (5-HT) by inhibitory feedback. The antagonistic properties of NAS-181 ((R)-(+)-2-[[[3-(Morpholinomethyl)-2H-chromen-8-ylloxy]methyl] morpholine methane sulfonate), a new selective antagonist for the rodent 5-HT_{1B} receptor, were determined by using an agonist-induced decrease of extracellular 5-HT. The 5-HT_{1B} receptor agonist CP93129 (0.030, 3 microm) applied by reversed microdialysis, dose-dependently reduced 5-HT levels in rat frontal cortex. The suppressant effect of CP93129 (0.1 microm) was smaller in the presence of fluvoxamine (3-10 microm), a 5-HT reuptake inhibitor. The effects of NAS-181 on CP93129 were compared with GR127935, a mixed 5-HT_{1B}/1D receptor antagonist, and SB224289, a 5-HT_{1B} receptor antagonist. Both in the presence and absence of fluvoxamine, the suppressant effect of CP93129 on extracellular 5-HT was attenuated by NAS-181 (1 microm) and GR127935 (10 microm), but not by SB224289 (1 microm). In the absence of fluvoxamine, GR127935, SB224289 and NAS-181 all reduced 5-HT levels, suggesting partial agonistic properties of these compounds. In conclusion, the results show that NAS-181 is a potent 5-HT_{1B} receptor antagonist.

L5 ANSWER 4 OF 4 MEDLINE ON STN
 ACCESSION NUMBER: 2003094519
 DOCUMENT NUMBER: PubMed ID: 12606921
 TITLE: The role of 5-HT_{1A/B} autoreceptors in the antinociceptive effect of systemic administration of acetaminophen.
 AUTHOR: Roca-Vinardell Aranzazu; Ortega-Alvaro Antonio; Gilbert-Ranola Juan; Mico Juan A
 CORPORATE SOURCE: Department of Neuroscience, Faculty of Medicine, University of Cadiz, Spain.
 SOURCE: Anesthesiology, (2003 Mar) Vol. 98, No. 3, pp. 741-7.
 Journal code: 1300217. ISSN: 0003-3022.
 United States
 Journal: Article; (JOURNAL ARTICLE)
 DOCUMENT TYPE: (RESEARCH SUPPORT, NON-U.S. GOV'T)
 LANGUAGE: English
 FILE SEGMENT: Abridged Index Medicus Journals; Priority Journals
 ENTRY MONTH: 200303
 ENTRY DATE: Entered STN: 28 Feb 2003
 Last Updated on STN: 28 Mar 2003
 Entered Medline: 27 Mar 2003

AB BACKGROUND: It has been proposed that serotonin participates in the central antinociceptive effect of acetaminophen. The serotonin activity in the brainstem is primarily under the control of 5-HT_{1A} somatodendritic receptors, although some data also suggest the involvement of 5-HT_{1B} receptors. In the presence of serotonin, the blockade of 5-HT_{1A/B} receptors at the level of the raphe nuclei leads to an increase in serotonin release in terminal areas, thus improving serotonin functions. This study examines the involvement of 5-HT_{1A/B} receptors in the antinociceptive effect of acetaminophen in mice. METHODS: The effects of acetaminophen (600 mg/kg intraperitoneal) followed by different doses of antagonists (WAY 100635 [0.2-0.8 mg/kg subcutaneous] and SB 216641 [0.2-0.8 mg/kg subcutaneous]) or agonists (8-OH-DPAT [0.25-1 mg/kg subcutaneous] and CP 93129 [0.125-0.5 mg/kg subcutaneous]) of 5-HT_{1A} and 5-HT_{1B} receptors, respectively, were determined in the hot-plate test in mice. RESULTS: Acetaminophen (300-800 mg/kg) showed a dose-dependent antinociceptive effect in the hot-plate test in mice. WAY

100635 (0.2-0.8 mg/kg; 5-HT_{1A} antagonist) induced an increase in the antinociceptive effect of 600 mg/kg acetaminophen, but this increase was not dose related. Conversely, 8-OH-DPAT (0.25-1 mg/kg; 5-HT_{1A} agonist) decreased the antinociceptive effect of acetaminophen. SB 216641 (0.2-0.8 mg/kg; 5-HT_{1B} antagonist) induced a dose-related increase in the antinociceptive effect of acetaminophen, and CP 93129 (0.25 mg/kg; 5-HT_{1B} agonist) significantly decreased the antinociceptive effect of acetaminophen. CONCLUSIONS: These results suggest that the combination of acetaminophen with compounds having 5-HT_{1A} and 5-HT_{1B} antagonist properties could be a new strategy to improve the analgesia of acetaminophen, thanks to its mild serotonergic properties.

=> s 2003 and d2 and (5HT1b or 5ht-1b or 5-HT1b) and (5ht2a or 5ht-2a or 5-HT2a)
 46652 2003
 26201 D2
 137 5HT1B
 2604 5HT
 13798 1B
 55 5HT-1B
 2352638 5 (5HT(W)1B)
 1858 HT1B
 1855 5-HT1B
 179 5HT2A
 2604 5HT
 23242 2A
 78 5HT-2A
 2352638 5 (5HT(W)2A)
 2592 HT2A
 2579 5-HT2A
 0 2003 AND D2 AND (5HT1B OR 5HT-1B OR 5-HT1B) AND (5HT2A OR 5HT-2A OR 5-HT2A)
 L6

=> s 2003/py and d2 and (5ht1b or 5ht-1b or 5-HT1b) and (5ht2a or 5ht-2a or 5-HT2a)
 569314 2003/PY
 (26030000-20039999/PY)
 26201 D2
 137 5HT1B
 2604 5HT
 13798 1B
 55 5HT-1B
 2352638 5 (5HT(W)1B)
 1858 HT1B
 1855 5-HT1B
 179 5HT2A
 2604 5HT
 23242 2A
 78 5HT-2A
 2352638 5 (5HT(W)2A)
 2592 HT2A
 2579 5-HT2A
 1 2003/PY AND D2 AND (5HT1B OR 5HT-1B OR 5-HT1B) AND (5HT2A OR 5HT-2A OR 5-HT2A)
 L7

=> d

L7 ANSWER 1 OF 1 MEDLINE on STN
 AN 2003157926 MEDLINE
 DN Pubmed ID: 12650952
 TI Investigating the role of dopaminergic and serotonergic candidate genes in
 obsessive-compulsive disorder.
 AU Hemmings Sian M J; Kinnear Craig J; Niehaus Dana J H; Moolman-Smook
 Johanna C; Lochner Christine; Knowles James A; Corfield Valerie A; Stein
 Dan J
 CS MRC/US Centre for Molecular and Cellular Biology, University of
 Stellenbosch, P.O. Box 19063, 7505, Tygerberg, South Africa...
 smjh@sun.ac.za
 SO Neuropsychopharmacology : the Journal of the European College of
 Neuropsychopharmacology, (2003 Mar) Vol. 13, No. 2, pp. 93-8.
 Journal code: 9111390. ISSN: 0924-977X.
 CY Netherlands
 DT (COMPARATIVE STUDY)
 Journal; Article; (JOURNAL ARTICLE)
 (RESEARCH SUPPORT, NON-U.S. GOV'T)
 LA English
 FS Priority Journals
 EM 200305
 ED Entered STN: 6 Apr 2003
 Last Updated on STN: 13 May 2003
 Entered Medline: 12 May 2003

=> d abs

L7 ANSWER 1 OF 1 MEDLINE on STN
 AB There is increasing evidence that the aetiology of obsessive-compulsive
 disorder (OCD) has a marked genetic component, although the precise
 mechanism of inheritance is unclear. Clinical and pharmacological studies
 have implicated the serotonergic and dopaminergic systems in disease
 pathogenesis. This study investigated the role of attractive candidate
 genes in the serotonergic and dopaminergic pathways in the development of
 OCD. The distribution of selected polymorphic variants in the serotonin
 receptor type 2A and 1b (5-HT(2A), 5-HT(1b)), dopamine transporter
 (DAT), dopamine receptor type 4 (DRD4) and monoamine-oxidase A (MAO-A)
 genes were analysed in 71 OCD cases and 129 control individuals in the
 genetically homogeneous Afrikaner population, by means of case-control
 association studies. Although no statistically significant genotypic or
 allelic associations were detected, the data yielded interesting
 preliminary results that warrant further discussion and investigation.

=> s d2 and (5ht1b or 5ht-1b or 5-ht1b) and (5ht2a or 5ht-2a or 5-ht2a)

26201 D2
 137 5HT1B
 2604 5HT
 13798 1B
 55 5HT-1B
 2352638 5 (5HT (W)1B)
 1858 HT1B
 1855 5-HT1B
 (5(W)HT1B)
 179 5HT2A
 2604 5HT
 23242 2A
 78 5HT-2A
 2352638 5 (5HT (W)2A)
 2592 HT2A
 2579 5-HT2A
 (5(W)HT2A)

L8 15 D2 AND (5HT1B OR 5HT-1B OR 5-HT1B) AND (5HT2A OR 5HT-2A OR 5-HT2
 A)

=> d 1-15 ibib abs hitstr
 'HITSTR' IS NOT A VALID FORMAT FOR FILE 'MEDLINE'

The following are valid formats:

The default display format is BIB.

ABS ---- AB
 ALL ---- AN, DN, TI, AU, CS, NC, SO, CM, CY, DT, LA, FS, NCT, OS,
 EM, ED, AB, ST, CT, NA, RN, CN, GEN
 BIB ---- AN, DN, TI, AU, CS, NC, SO, CY, DT, LA, FS, NCT, OS, EM, ED
 CBIB ---- AN, DN, TI, AU, CS, NC, SO, CY, DT, LA, FS, NCT, OS, EM, ED
 DALL ---- ALL, delimited for post processing
 IABS ---- ABS, with a text label
 IALL ---- ALL, indented with text labels
 IBIB ---- BIB, indented with text labels
 IND ---- ST, CT, NA, RN, CN, GEN
 TRIAL -- TI, ST, CT, NA, RN, CN, GEN
 (SAM, TRI, FREE)
 HIT ---- All fields containing hit terms
 HITIND - IND
 KWIC --- All hit terms plus 20 words on either side
 OCC ---- List of display fields containing hit terms

Hit terms will be highlighted in all available fields except CM and
 PY.

To display a particular field or fields, enter the display field
 codes. For a list of display field codes, enter 'HELP DFIELDS'
 at an arrow prompt (=>). Examples of formats include: 'BIB';
 'AB'; 'SO,ST'. You may specify the format fields in any order,
 and the information will be displayed in the same order as the
 format specification.

The same formats (except for HIT, HITIND, KWIC, and OCC) may be
 used with the DISPLAY ACC command to display the record for a
 specified Accession Number.

ENTER DISPLAY FORMAT (BIB):end

=> d 1-15 ibib abs

L8 ANSWER 1 OF 15 MEDLINE on STN
 ACCESSION NUMBER: 2006514916
 DOCUMENT NUMBER: Pubmed ID: 16820177
 TITLE: Quantitative mapping shows that serotonin rather than
 dopamine receptor mRNA expressions are affected after
 repeated intermittent administration of MDMA in rat brain.
 AUTHOR: Kindlundh-Hogberg Anna M S; Svenningsson Per; Schioth Helgi
 B
 CORPORATE SOURCE: Department of Physiology and Pharmacology, Karolinska
 Institutet, Stockholm, Sweden.. anna.kindlundh@neuro.uu.se
 SOURCE: Neuropsychopharmacology, (2006 Sep) Vol. 31, No. 4, pp. 838-47.
 PUB. COUNTRY: Electronic Publication: 2006-07-03
 DOCUMENT TYPE: Journal code: 0236217. ISSN: 0028-3908.
 (COMPARATIVE STUDY)
 Journal; Article; (JOURNAL ARTICLE)
 (RESEARCH SUPPORT, NON-U.S. GOV'T)
 LANGUAGE: English
 FILE SEGMENT: Priority Journals
 ENTRY MONTH: 200611
 ENTRY DATE: Entered STN: 30 Aug 2006

Last Updated on STN: 7 Nov 2006

Entered Medline: 6 Nov 2006

AB Ecstasy, (+/-)-3,4-methylenedioxymetamphetamine (MDMA), is a popular recreational drug among young people. The present study aims to mimic MDMA intake among adolescents at dance clubs, taking repeated doses in the same evening on an intermittent basis. Male Sprague-Dawley rats received either 3x1 or 3x5 mg/kg/day (3 h apart) every seventh day during 4 weeks. We used real-time RT-PCR to determine the gene expression of serotonin 5HT1A, 5HT1B, 5HT2A, 5HT2C, 5HT3, 5HT6 receptors and the dopamine D1, D2, D3 receptors in seven brain nuclei. The highest dose of MDMA extensively increased the 5HT1B-receptor mRNA in the cortex, caudate putamen, nucleus accumbens, and hypothalamus. The 5HT2A-receptor mRNA was reduced at the highest MDMA dose in the cortex. The 5HT2C mRNA was significantly increased in a dose-dependent manner in the cortex and the hypothalamus, as well as the 5HT3-receptor mRNA was in the hypothalamus. The 5HT6 mRNA level was increased in the forebrain cortex and the amygdala. Dopamine receptor mRNAs were only affected in the hypothalamus. In conclusion, this study provides evidence for a unique implication of serotonin rather than dopamine receptor mRNA levels, in response to repeated intermittent MDMA administration. We therefore suggest that serotonin regulated functions also primarily underlie repeated MDMA intake at rave parties.

L8 ANSWER 2 OF 15 MEDLINE on STN
2003157926

ACCESSION NUMBER: PubMed ID: 12650952

DOCUMENT NUMBER: Investigating the role of dopaminergic and serotonergic candidate genes in obsessive-compulsive disorder.

AUTHOR: Hemmings Sian M J; Kinnear Craig J; Niehaus Dana J H; Moolman-Smook Johanna C; Lochner Christine; Knowles James A; Corfield Valerie A; Stein Dan J

CORPORATE SOURCE: MRC/JUS Centre for Molecular and Cellular Biology, University of Stellenbosch, P.O. Box 19063, 7505, Tygerberg, South Africa. smjhesus.ac.za
SOURCE: European neuropsychopharmacology : the journal of the European College of Neuropsychopharmacology, (2003 Mar) Vol. 13, No. 2, pp. 93-8.
Journal code: 9111390. ISSN: 0924-977X.
Netherlands

PUB. COUNTRY: (COMPARATIVE STUDY)
DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)

LANGUAGE: (RESEARCH SUPPORT, NON-U.S. GOV'T)

FILE SEGMENT: English

ENTRY MONTH: Priority Journals

ENTRY DATE: 200305

Entered STN: 6 Apr 2003

Last Updated on STN: 13 May 2003

Entered Medline: 12 May 2003

AB There is increasing evidence that the aetiology of obsessive-compulsive disorder (OCD) has a marked genetic component, although the precise mechanism of inheritance is unclear. Clinical and pharmacological studies have implicated the serotonergic and dopaminergic systems in disease pathogenesis. This study investigated the role of attractive candidate genes in the serotonergic and dopaminergic pathways in the development of OCD. The distribution of selected polymorphic variants in the serotonin receptor type 2A and 1D beta (5-HT(2A), 5-HT(1D beta)), dopamine transporter receptor type 4 (DRD4), and monoamine-oxidase A (MAO-A) genes were analysed in 71 OCD cases and 129 control individuals in the genetically homogeneous Afrikaner population, by means of case-control association studies. Although no statistically significant genotypic or allelic associations were detected, the data yielded interesting preliminary results that warrant further discussion and investigation.

L8 ANSWER 3 OF 15 MEDLINE on STN
2001300306

ACCESSION NUMBER: MEDLINE

DOCUMENT NUMBER:
TITLE:

AUTHOR:

CORPORATE SOURCE:

SOURCE:

PUB. COUNTRY:

DOCUMENT TYPE:

LANGUAGE:

FILE SEGMENT:

ENTRY MONTH:

ENTRY DATE:

AB It is becoming increasingly clear that environmental stimuli play a critical role in the maintenance of drug taking behaviour. This has led to investigations into the neural mechanisms by which environmental stimuli can come to control behaviour using paradigms such as conditioned reinforcement. The majority of this work has involved the use of food-paired conditioned stimulus rodent paradigms. Relatively few studies have attempted to investigate the neuropharmacology of behaviour maintained by presentation of a stimulus paired with ethanol drinking. Several lines of research support an important role for brain serotonin (5-HT) neurotransmitter systems in the control of alcohol drinking behaviour. The aim of the present study was, initially, to establish a procedure in which rats respond for an ethanol-paired conditioned stimulus, and second, to study the effects of a range of serotonergic compounds previously shown to be effective in reducing oral ethanol self-administration, on responding for this conditioned stimulus. Results showed that the 5-HT releaser d-fenfluramine, the selective serotonin reuptake inhibitor fluoxetine, the 5-HT1A receptor agonist 8-hydroxy-2(di-n-propylamino)tetralin, the partial 5-HT1A receptor agonist 1-(3-trifluoromethylphenyl)piperazine, but not the 5-HT2A/5-HT2C receptor agonist 1-(2,5-dimethoxy-4-iodophenylaminopropane)-2, selectively reduced responding on a lever leading to presentation of an ethanol paired conditioned stimulus. In addition the non-specific D1/D2 dopamine receptor antagonist haloperidol was active in this paradigm. Results are consistent with involvement of the dopaminergic and 5-HT systems, in particular activation of 5-HT1A and 5-HT1B receptor subtypes, in mediation of the conditioned or secondary reinforcing properties of ethanol.

L8 ANSWER 4 OF 15 MEDLINE on STN

ACCESSION NUMBER: 2001170940 MEDLINE

DOCUMENT NUMBER: PubMed ID: 11271402

TITLE: Receptor-mediated regulation of serotonin output in the rat dorsal raphe nucleus: effects of risperidone.

AUTHOR: Hertel P; Lindblom N; Nomikos G G; Svensson T H

CORPORATE SOURCE: Department of Physiology and Pharmacology, Karolinska Institutet, Stockholm, Sweden.

SOURCE: Psychopharmacology, (2001 Jan) Vol. 153, No. 3, pp. 307-14. Journal code: 7608025. ISSN: 0033-3158.

PUB. COUNTRY: Germany; Germany, Federal Republic of

DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)

LANGUAGE: English

FILE SEGMENT: Priority Journals

ENTRY MONTH: 200105

ENTRY DATE: Entered STN: 21 May 2001

PubMed ID: 11198050

Manipulation of operant responding for an ethanol-paired conditioned stimulus in the rat by pharmacological alteration of the serotonergic system.

Wilson A W; Costall B; Neill J C

Postgraduate Studies in Pharmacology, School of Pharmacy, University of Bradford, West Yorkshire, UK...

alex.wilson@bphrd.com

Journal of psychopharmacology (Oxford, England), (2000) Vol. 14, No. 4, pp. 340-6.

Journal code: 8907828. ISSN: 0269-8811.

United States

Journal; Article; (JOURNAL ARTICLE)

English

Priority Journals

200105

Entered STN: 4 Jun 2001

Last Updated on STN: 4 Jun 2001

Entered Medline: 31 May 2001

Last Updated on STN: 21 May 2001

Entered Medline: 17 May 2001

AB OBJECTIVES: The present study was undertaken to characterize the regulation of serotonin (5-HT) efflux and neuronal activity in the dorsal raphe nucleus (DRN) as well as to examine the potential ability of the antipsychotic drug risperidone to interfere with these mechanisms. **METHODS AND RESULTS:** By using microdialysis in freely moving rats, it was found that administration of the alpha2-adrenoceptor antagonist idazoxan (0.25 mg/kg, SC), the 5-HT1B/D receptor antagonist GR 127,935 or 127,935 (1.0 mg/kg, SC) and risperidone (0.6 or 2.0 mg/kg, SC) increased 5-HT output in the DRN. Local DRN perfusion with GR 127,935 or risperidone via reversed dialysis (100 or 10-100 microm, respectively) enhanced 5-HT efflux in this area, whereas idazoxan (10-100 microm) failed to affect this parameter. Both systemic administration and reversed DRN dialysis of the D2/3 and 5-HT2A receptor antagonist GR 127,935 or antagonists raclopride (2.0 mg/kg, SC or 10-100 microm) and MDL 100,907 (1.0 mg/kg, SC or 10-100 microm), respectively, were without effect. Intrathecal dialysis of the 5-HT1B/D receptor agonist CP 135,807 (0.2 microm) decreased the efflux of 5-HT in the DRN, an effect which was antagonized by co-administration of either GR 127,935 or risperidone (10 and 3.3 microm, respectively). By using single-cell recording, it was found that administration of GR 127,935 (50-400 microg/kg, IV) decreased, whereas CP 135,807 (2.5-20 microg/kg, IV) increased firing of 5-HT cells in the DRN. **CONCLUSIONS:** Our findings suggest a regulatory role of local 5-HT1B/D receptors on 5-HT efflux as well as cell firing in the DRN and indicate that risperidone may interfere with the regulation of 5-HT availability in this area primarily via blockade of 5-HT1D receptors.

L8 ANSWER 5 OF 15 MEDLINE on STN

ACCESSION NUMBER: 1999371966 MEDLINE

DOCUMENT NUMBER: PubMed ID: 10443019

TITLE: 2-[4-[3-(4-aryl/heteroaryl-1-piperazinyl)propoxy]phenyl]-2H-benzotriazoles and their N-oxides as ligands for serotonin and dopamine receptors.

AUTHOR: Sparatore A; Gogean M; Cagnotto A; Sparatore F

CORPORATE SOURCE: Istituto di Chimica Farmaceutica e Tossicologica,

Università di Milano, Italy.. sparatore@iuc.chemi.unimi.it

SOURCE: Farmaco (Società chimica italiana : 1989), (1999 Jun 30)

Vol. 54, No. 6, pp. 402-10.

JOURNAL CODE: 8912641. ISSN: 0014-827X.

PUB. COUNTRY: Italy

DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)

LANGUAGE: English

ENTRY MONTH: Priority Journals

ENTRY DATE: 199911

ENTRY DATE: Entered STN: 11 Jan 2000

ENTRY DATE: Last Updated on STN: 11 Jan 2000

ENTRY DATE: Entered Medline: 10 Nov 1999

AB A small set of 2-[4-[3-(4-aryl/heteroaryl-piperazinyl)propoxy]phenyl]-2H-benzo tri azoles and corresponding N-oxides were prepared. The synthesized compounds were able to bind on some serotonin (5-HT1A, 5-HT2A) and dopamine (D2, D3) receptors, while displaying poor or no affinity for 5-HT1B, 5-HT2C, 5-HT3, and 5-HT4 subtypes. The strong contribution of the N-oxide function for the binding on 5-HT1A, D2 and D3 receptors is noteworthy. For 2-[4-[3-(4-(2-methoxyphenyl)-1-piperazinyl)propoxy]phenyl]-2H-benzotriazole-1-oxide (4b), the binding constants (K1) were 11.9 (5-HT1A) and 10.5 nM (D3). In a general pharmacological screening, the 2-[4-[3-(4-phenyl)-1-piperazinyl]propoxy]phenyl]-2H-benzotriazole (3a) exhibited only very weak activities, with the exception of protecting mice from cyanide-induced hypoxia.

L8 ANSWER 6 OF 15 MEDLINE on STN

ACCESSION NUMBER: 1998389363 MEDLINE

DOCUMENT NUMBER: PubMed ID: 9723970

TITLE: The role of alpha2-adrenoceptor antagonism in the anti-cateleptic properties of the atypical neuroleptic agent, clozapine, in the rat.

AUTHOR: Kalkman H O; Neumann V; Hoyer D; Tricklebank M D

CORPORATE SOURCE: Nervous Systems Research, Novartis Pharma AG, Basel,

SOURCE: British Journal of pharmacology, (1998 Aug) Vol. 124, No. 7, pp. 1550-6.

JOURNAL CODE: 7502536. ISSN: 0007-1188.

PUB. COUNTRY: ENGLAND: United Kingdom

DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)

LANGUAGE: English

ENTRY MONTH: Priority Journals

ENTRY DATE: 199811

ENTRY DATE: Entered STN: 6 Jan 1999

ENTRY DATE: Last Updated on STN: 6 Jan 1999

ENTRY DATE: Entered Medline: 13 Nov 1998

AB 1. The mechanism underlying the anticeleptic properties of the atypical neuroleptic agent, clozapine, has been investigated in the rat. 2. The close structural analogues of clozapine, loxapine (0.1 mg kg(-1) s.c.) and iso-clozapine (1 and 3 mg kg(-1) s.c.) induced catalepsy in rats. In contrast, clozapine and the regio-isomer of loxapine, iso-loxapine (up to 10 mg kg(-1) s.c.) did not produce catalepsy, but at a dose of 1 mg kg(-1) significantly inhibited catalepsy induced by loxapine (0.3 mg kg(-1) s.c.). 3. Radioligand binding assays showed that cataleptic potential was most clearly predicted by the D2/5-HT1A, D2/5-HT1B/1D and D2/alpha2-receptor affinity (K0) ratios: i.e. 30-100-fold higher ratios were calculated for loxapine and iso-clozapine, whereas the ratios were less than 1 for clozapine and iso-loxapine. The ratios of affinities for D2 to 5-HT2A, 5-HT2C or D1 did not reflect the grouping of cataleptic and non-cataleptic compounds. 4. Co-treatment with the alpha2-adrenoceptor antagonists, yohimbine (1-10 mg kg(-1) s.c.), RX 821002 (1-10 mg kg(-1) s.c.) and MK-912 (0.3 and 1 mg kg(-1) s.c.) dose-dependently inhibited the cataleptic response to loxapine (0.3 mg kg(-1) s.c.). Yohimbine (1-10 mg kg(-1) s.c.) also dose-dependently inhibited the cataleptic response to haloperidol (0.3 mg kg(-1) s.c.). The alpha2-adrenoceptor antagonists had no effect per se. 5. Neither yohimbine (10 mg kg(-1)) nor RX821002 (3 mg kg(-1)) altered the cataleptic response to the D1 receptor antagonist, SCH 23390 (1 mg kg(-1) s.c.), while, like clozapine, both compounds abolished the response to the 5-HT2A receptor antagonist, MDL 100,151 (3 mg kg(-1) s.c.). 6. The present data strongly implicate alpha2-adrenoceptor blockade in the anticeleptic properties of clozapine and suggest that its lack of extrapyramidal side effects in the clinic may also be a consequence of this property.

L8 ANSWER 7 OF 15 MEDLINE on STN

ACCESSION NUMBER: 1998316312 MEDLINE

DOCUMENT NUMBER: PubMed ID: 9651340

TITLE: Cross-talk between 5-hydroxytryptamine receptors in a serotonergic cell line. Involvement of arachidonic acid metabolism.

AUTHOR: Tournais C; Mutel V; Manivet P; Launay J M; Kellermann O

CORPORATE SOURCE: Différenciation Cellulaire, CNRS URA 1960, Institut

Pasteur, 25 rue du Dr. Roux, 75724 Paris Cedex 15, France.

SOURCE: The Journal of biological chemistry, (1998 Jul 10) Vol. 273, No. 28, pp. 17498-503.

JOURNAL CODE: 2985121R. ISSN: 0021-9258.

PUB. COUNTRY: United States

DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)

LANGUAGE: (RESEARCH SUPPORT, NON-U.S. GOV'T)

LANGUAGE: English

FILE SEGMENT: 199808
ENTRY MONTH: 199808
ENTRY DATE: 17 Aug 1998
Last Updated on STN: 17 Aug 1998
Entered Medline: 6 Aug 1998

AB The study of signaling cascades and of functional interactions between cell populations remains an arduous task. We took advantage of a serotonergic cell line to elucidate cross-talks between 5-HT receptors and to demonstrate the involvement of two 5-HT2 receptor subtypes in the regulation of 5-HT1B/1D function. The inducible 1C11 cell line has the unique property of acquiring within 4 days a complete serotonergic phenotype (1C11+ cells), including three 5-HT receptors. 5-HT1B/1D and 5-HT2B receptors are expressed since day 2 of the serotonergic differentiation while 5-HT2A receptors are induced at day 4. We first established that 5-HT2B receptors are coupled with the phospholipase A2 (PLA2)-mediated release of arachidonic acid (AA) and that the activation of 5-HT2B receptors in 1C11+ d2 cells inhibits the 5-HT1B/1D receptor function via a cyclooxygenase-dependent AA metabolite. At day 4, this 5-HT2B-mediated inhibition of the 5-HT1B/1D function can be blocked upon concomitant 5-HT2A activation although a 5-HT2B/PLA2 positive coupling was evidenced. This suggests the existence in 1C11+d4 cells of pathway(s) for 5-HT2A receptors, distinct from PLC and PLA2. Finally, this study reveals the antagonistic roles of 5-HT2A and 5-HT2B receptors in regulating the function of 5-HT1B/1D, a receptor involved in neuropsychiatric disorders and migraine pathogenesis.

L8 ANSWER 8 OF 15 MEDLINE on STN
ACCESSION NUMBER: 1998230395
DOCUMENT NUMBER: Pubmed ID: 9570468

TITLE: Cataleptogenic effect of subtype selective 5-HT receptor antagonists in the rat.

AUTHOR: Kalkman H O; Neumann V; Nozulak J; Tricklebank M D
CORPORATE SOURCE: Nervous System Research, Novartis Pharma Inc., Basel, Switzerland.
SOURCE: European Journal of pharmacology, (1998 Feb 19) Vol. 343, No. 2-3, pp. 201-7.
Journal code: 1254354. ISSN: 0014-2999.

PUB. COUNTRY: Netherlands
DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)
LANGUAGE: English
FILE SEGMENT: Priority Journals
ENTRY MONTH: 199806
ENTRY DATE: 18 Jun 1998
Last Updated on STN: 18 Jun 1998
Entered Medline: 9 Jun 1998

AB 5-HT receptor antagonists with selectivity for 5-HT1A WAY-100635 (N-[2-[4-(2-methoxyphenyl)-1-piperazinyl]ethyl]-N-(2-pyridinyl)cyclohexanecarboxamide), 5-HT1B GR 127935 (N-(methoxy-3-(4-methyl-1-piperazinyl)phenyl)-2'-methyl-4'-(5-methyl-1,2,4-oxadiazol-3-yl)[1,1-biphenyl]-4-carboxamide x HCl), 5-HT2C SB 200646A (N-(1-methyl-5-indolyl)-N'-(3-pyridyl)urea x HCl) and 5-HT2A (ketanserin, fanserin and MDL 100,151 ((+/-)-alpha-(2,3-dimethoxyphenyl)-1-[2-(4-fluorophenyl)ethyl]-4-piperidinemethanol) receptors were tested for cataleptogenic responses in rats. WAY-100635 (0.1-3 mg/kg, s.c.), ketanserin (0.1-3 mg/kg, s.c.), MDL 100,151 (0.3-3 mg/kg, s.c.) and fanserin (RP 62203; 3 mg/kg, s.c.) induced a significant catalepsy. GR 127935 (1 mg/kg, s.c.), SB 200646A (without effect per se at 10 mg/kg, s.c.) and MDL 100,151 (0.3 mg/kg, s.c.) did not inhibit the cataleptic response to the dopamine D2 receptor antagonist, loxapine (0.3 mg/kg, s.c.). Catalepsy induced by MDL 100,151 (3 mg/kg) was blocked by co-treatment with clozapine, but not by SB

200646A (both at 10 mg/kg, s.c.). Although clozapine displays significant affinity to 5-HT1A, 5-HT1B, 5-HT2A and 5-HT2C receptors, the present results suggest that blockade of these receptors is not responsible for clozapine's anticonvulsant activity.

L8 ANSWER 9 OF 15 MEDLINE on STN
ACCESSION NUMBER: 1998149351
DOCUMENT NUMBER: Pubmed ID: 9489707

TITLE: Serotonin neural adaptations to ontogenetic loss of dopamine neurons in rat brain.

AUTHOR: Kostzewa R M; Reader T A; Descarries L
CORPORATE SOURCE: Department of Pharmacology, James H. Quillen College of Medicine, East Tennessee State University, Johnson City, USA.

SOURCE: Journal of neurochemistry, (1998 Mar) Vol. 70, No. 3, pp. 889-98. Ref: 94
Journal code: 2985190R. ISSN: 0022-3042.

PUB. COUNTRY: United States
DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)
LANGUAGE: English
FILE SEGMENT: General Review; (REVIEW)
ENTRY MONTH: Priority Journals
ENTRY DATE: 199803

AB In rat, the neonatal destruction of nigrostriatal dopamine (DA) neurons by intracerebral administration of 6-hydroxydopamine entails dramatic changes in serotonin (5-HT) hyperinnervation of the adult neostriatum, associated with increases in density of various 5-HT receptor subtypes and enhanced neuronal responsiveness to the iontophoretic application of 5-HT and its 5-HT1B/2C and 5-HT2A/2C receptor agonists, m-chlorophenylpiperazine and iodoamethoxyphenylaminepropane. The topographical distribution of these changes is consistent with up-regulation and/or increased production and transport of 5-HT1B and 5-HT2A receptors by the neostriatal projection neurons, as confirmed for the 5-HT2A receptor in a recent in situ hybridization study. It is interesting that this study has also shown that increases in both 5-HT2A binding and mRNA level were abolished by chronic pretreatment with the DA agonists, apomorphine and SKF 38393, suggesting a regulatory influence of DA in the expression of this 5-HT receptor. D1 receptor binding is known to be slightly reduced in the rostral neostriatum of these rats, a down-regulation apparently imputable to a reduced rate of synthesis of the receptor. In contrast, D2 receptor binding is increased throughout the DA-denervated and 5-HT-hyperinnervated neostriatum, perhaps due to some posttranscriptional modifications. Stereotyped and motor behaviors induced by systemic treatment with D1 and D2 agonists are markedly enhanced in these rats (behavioral supersensitivity), although priming is commonly required to unmask a latent D1 supersensitivity. In the case of oral activity, however, overt behavioral supersensitivity is induced by D1 as well as D2 agonists. Moreover, there is overt supersensitivity of oral activity in response to the 5-HT receptor agonist m-chlorophenylpiperazine, which is presumably imputable to 5-HT2C receptors and may be demonstrated even in the absence of supersensitivity to D1 receptor agonist. 5-HT adaptations, therefore, seem to play a role not only in the abnormal spontaneous behavior, but also in the behavioral supersensitivity to 5-HT as well as DA receptor agonists in these rats.

L8 ANSWER 10 OF 15 MEDLINE on STN
ACCESSION NUMBER: 97021512
DOCUMENT NUMBER: Pubmed ID: 8867872

TITLE: Evidence that m-chlorophenylpiperazine-induced hyperthermia in rats is mediated by stimulation of 5-HT2C receptors.

AUTHOR: Mazzola-Pomietto P; Aulakh C S; Wozniak K M; Murphy D L
CORPORATE SOURCE: Laboratory of Clinical Science, National Institute of Mental Health, National Institutes of Health, Bethesda, MD 20892-1264, USA.

SOURCE: Psychopharmacology, (1996 Feb) Vol. 123, No. 4, pp. 333-9. Journal code: 7608025. ISSN: 0033-3158.

PUB. COUNTRY: GERMANY: Germany, Federal Republic of
DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)
LANGUAGE: English
FILE SEGMENT: Priority Journals
ENTRY MONTH: 199611
ENTRY DATE: Entered STN: 19 Dec 1996
 Last Updated on STN: 19 Dec 1996
 Entered Medline: 15 Nov 1996

AB Intraperitoneal administration of m-chlorophenylpiperazine (m-CPP) to Wistar rats produced hyperthermia with a peak effect at 30 min. Pretreatment with low doses of metergoline (5-HT1/5-HT2 antagonist), mesulergine and mianserin (5-HT2C/5-HT2A antagonists) blocked m-CPP-induced hyperthermia. Pretreatment with propranolol (beta-adrenergic receptor antagonist that also has binding affinity for 5-HT1A, 5-HT1B and 5-HT2B sites), yohimbine (alpha 2-noradrenergic antagonist that also has binding affinity for 5-HT2B sites), MDL-72222 or ondansetron (5-HT3 antagonists) did not attenuate m-CPP-induced hyperthermia. Only high doses of ketanserin, LY-53857 and ritanserin (5-HT2A/5-HT2C antagonists) as well as spiperone (5-HT1A/5-HT2A/D2 antagonist) attenuated m-CPP-induced hyperthermia. Daily administration of m-CPP produced complete tolerance to its hyperthermic effect by day 5. However, there was no cross-tolerance to 1-(2,5-dimethoxy-4-iodophenyl)-2-aminopropane (DOI), a 5-HT2A agonist that also has high affinity for 5-HT2C receptors)-induced hyperthermia. m-CPP-induced increases in temperature were found to be significantly less in the Fawn-Hooded (FH) rat strain as compared to the Wistar rat strain; in prior studies, FH rats have been found to be subsensitive to other hyperthermic agents. These findings suggest that m-CPP-induced hyperthermia in rats is mediated by selective stimulation of 5-HT2C receptors.

L8 ANSWER 11 OF 15 MEDLINE on STN
ACCESSION NUMBER: 95316519 MEDLINE
DOCUMENT NUMBER: PubMed ID: 7796132
TITLE: Short and long-term changes in dopamine and serotonin receptor binding sites in amphetamine-sensitized rats: a quantitative autoradiographic study.
 Bonhomme N; Cador M; Stinus L; Le Moal M; Spampinato U
AUTHOR: INSERM U. 259, Université de Bordeaux II, France.
CORPORATE SOURCE: Brain research, (1995 Mar 27) Vol. 675, No. 1-2, pp. 215-23.
SOURCE: Journal code: 0045503. ISSN: 0006-8993.

PUB. COUNTRY: Netherlands
DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)
LANGUAGE: (RESEARCH SUPPORT, NON-U.S. GOV'T)
FILE SEGMENT: English
ENTRY MONTH: Priority Journals
ENTRY DATE: Entered STN: 17 Aug 1995
 Last Updated on STN: 17 Aug 1995
 Entered Medline: 3 Aug 1995

AB The biochemical changes in DA and 5HT systems were investigated in amphetamine (AMPH)-sensitized rats, 1 and 15 days after cessation of treatment (5 mg/kg AMPH, i.p., twice a day for 6 days). At both times, AMPH-treated rats exhibited behavioral sensitization, as revealed by an enhancement of the stereotypic response to a challenge dose of 2 mg/kg, ip. AMPH. Basal dopamine (DA) and serotonin (5-HT) metabolism was not significantly modified in different brain areas of AMPH-sensitized rats.

Quantitative autoradiographic analysis of DA and serotonin 5-HT receptor subtypes was performed in the following brain regions: medial prefrontal cortex, nucleus accumbens, striatum, substantia nigra, ventral tegmental area, dorsal and medial raphe nuclei. A significant increase of [3H]SCH 23390 binding to D1 DA receptors was observed in the substantia nigra pars reticulata 1 day but not 15 days after the cessation of AMPH treatment, whereas [3H]8-OH-DPAT binding to 5-HT1A sites was found to be significantly enhanced in the dorsal raphe nucleus at both time points. No change in D2 DA nor in 5-HT1B or 5-HT2A receptors was found in any of the brain structures examined at either time point. The obtained results suggest that DA and 5-HT systems are differently and time-dependently involved in AMPH-induced behavioral sensitization.

L8 ANSWER 12 OF 15 MEDLINE on STN
ACCESSION NUMBER: 95273535 MEDLINE
DOCUMENT NUMBER: PubMed ID: 7753967
TITLE: Evidence that 1-(2,5-dimethoxy-4-iodophenyl)-2-aminopropane (DOI)-induced hyperthermia in rats is mediated by stimulation of 5-HT2A receptors.
 Mazzola-Pomietto P; Aulakh C S; Wozniak K M; Hill J L; Murphy D L
AUTHOR: Laboratory of Clinical Science, National Institute of Mental Health, National Institutes of Health, Bethesda, MD 20892, USA.
CORPORATE SOURCE: Psychopharmacology, (1995 Jan) Vol. 117, No. 2, pp. 193-9. Journal code: 7608025. ISSN: 0033-3158.
SOURCE: GERMANY: Germany, Federal Republic of
PUB. COUNTRY: Journal; Article; (JOURNAL ARTICLE)
DOCUMENT TYPE: English
LANGUAGE: Priority Journals
FILE SEGMENT: 199506
ENTRY MONTH: Entered STN: 29 Jun 1995
ENTRY DATE: Last Updated on STN: 29 Jun 1995
 Entered Medline: 22 Jun 1995

AB The effects of various 5-HT receptor subtype-selective antagonists were studied on phenylisopropylamine hallucinogen 1-(2,5-dimethoxy-4-iodophenyl)-2-aminopropane (DOI)-induced hyperthermia in Wistar rats, in an attempt to characterize the 5-HT receptor subtype mediating DOI-induced hyperthermia. Intraperitoneal administration of DOI to rats produced hyperthermia with a peak effect at 60 min. Pretreatment with propranolol (beta-adrenoceptor antagonist that also has binding affinity for 5-HT1A, 5-HT1B and 5-HT2C sites), MDL-72222 or ondansetron (5-HT3 antagonists) did not attenuate DOI-induced hyperthermia. In contrast, pretreatment with metergoline (5-HT1/5-HT2 antagonist), ketanserin, LY53857, mesulergine, mianserin and ritanserin (5-HT2C/5-HT2A antagonists), as well as spiperone (5-HT1A/5-HT2A/D2 antagonist), significantly attenuated DOI-induced hyperthermia. Furthermore, daily administration of DOI (2.5 mg/kg per day) for 17 days did not produce either tolerance to its hyperthermic effect or modify m-CPP-induced hyperthermia in rats. These findings suggest that DOI-induced hyperthermia in rats is mediated by stimulation of 5-HT2A receptors.

L8 ANSWER 13 OF 15 MEDLINE on STN
ACCESSION NUMBER: 95055070 MEDLINE
DOCUMENT NUMBER: PubMed ID: 7965707
TITLE: Role of various 5-HT receptor subtypes in mediating neuroendocrine effects of 1-(2,5-dimethoxy-4-methylphenyl)-2-aminopropane (DOM) in rats.
 Aulakh C S; Mazzola-Pomietto P; Hill J L; Murphy D L
AUTHOR: Laboratory of Clinical Science, National Institute of Mental Health, Bethesda, Maryland
CORPORATE SOURCE: The Journal of pharmacology and experimental therapeutics, (1994 Oct) Vol. 271, No. 1, pp. 143-8.
SOURCE:

Journal code: 0376362. ISSN: 0022-3565.

PUB. COUNTRY:
DOCUMENT TYPE:
LANGUAGE:
FILE SEGMENT:
ENTRY MONTH:
ENTRY DATE:

United States
Journal; Article; (JOURNAL ARTICLE)
English
Priority Journals
199411
Entered STN: 10 Jan 1995
Last Updated on STN: 10 Jan 1995
Entered Medline: 25 Nov 1994

AB The phenylisopropylamine hallucinogen 1-(2,5-dimethoxy-4-methylphenyl)-2-aminopropane (DOM) produced dose-related increases in plasma concentrations of prolactin, adrenocorticotrophic hormone (ACTH) and corticosterone but not growth hormone in rats. Pretreatment with metergoline (serotonin, 5-HT₁/5-HT₂ antagonist), ritanserin and mianserin (5-HT_{2A}/5-HT_{2C} antagonists) significantly attenuated DOM-induced increases in prolactin, ACTH and corticosterone, whereas metergoline (5-HT_{2A}/5-HT_{2C} antagonist) pretreatment significantly attenuated DOM-induced increases in plasma prolactin and ACTH but not corticosterone. Pretreatment with propranolol (beta adrenoceptor antagonist that also has high binding affinity for 5-HT_{1A}, 5-HT_{1B} and 5-HT_{2C} sites), MDL-72222 and ondansetron (5-HT₃ antagonists) attenuated DOM's effect on plasma prolactin, but did not attenuate DOM-induced increases in either ACTH or corticosterone. On the other hand, spiperone (5-HT_{1A}/5-HT_{2A}/D₂ antagonist) pretreatment significantly attenuated DOM-induced increases in ACTH but not corticosterone. These findings demonstrate involvement of 5-HT_{2A}/5-HT_{2C} and 5-HT₃ receptors in mediating DOM-induced increases in plasma prolactin, whereas DOM-induced increases in ACTH appear to be mediated by stimulation of 5-HT_{2A} receptors. DOM-induced corticosterone secretion appears to be mediated by stimulation of 5-HT_{2A} and/or 5-HT_{2C} receptors. DOM does not affect growth hormone secretion in rats.

L8 ANSWER 14 OF 15

MEDLINE on STN

ACCESSION NUMBER:
DOCUMENT NUMBER:
TITLE:

94308931
Pubmed ID: 8035308
Evidence that 1-(2,5-dimethoxy-4-methylphenyl)-2-aminopropane-induced hypophagia and hyperthermia in rats is mediated by serotonin-2A receptors.

AUTHOR:

Aulakh C S; Mazzola-Pomietto P; Wozniak K M; Hill J L; Murphy D L

CORPORATE SOURCE:

Laboratory of Clinical Science, National Institute of Mental Health, Bethesda, Maryland.
The Journal of pharmacology and experimental therapeutics, (1994 Jul) Vol. 270, No. 1, pp. 127-32.
Journal code: 0376362. ISSN: 0022-3565.

SOURCE:

PUB. COUNTRY:
DOCUMENT TYPE:
LANGUAGE:
FILE SEGMENT:
ENTRY MONTH:
ENTRY DATE:

United States
Journal; Article; (JOURNAL ARTICLE)
English
Priority Journals
199408
Entered STN: 25 Aug 1994
Last Updated on STN: 25 Aug 1994
Entered Medline: 15 Aug 1994

AB The administration of various doses of the phenylisopropylamine hallucinogen 1-(2,5-dimethoxy-4-methylphenyl)-2-aminopropane (DOM) to rats produced dose-related decreases in 1-hr food intake in a food-restricted paradigm and in locomotor activity. DOM also produced dose-related increases in temperature. Pretreatment with propranolol (a beta adrenoceptor antagonist that also has high binding affinity for serotonin (5-HT) 5-HT_{1A}, 5-HT_{1B} and 5-HT_{2C} sites), metergoline or ondansetron (5-HT₃ antagonists) did not attenuate either DOM-induced hypophagia or hyperthermia. In contrast, pretreatment with metergoline (a 5-HT₁/5-HT₂ antagonist) and ritanserin (a 5-HT_{2A}/5-HT_{2C} antagonist) significantly attenuated both DOM-induced hypophagia

and hyperthermia. However, pretreatment with mesulergine (a 5-HT_{2C}/5-HT_{2A} antagonist) significantly attenuated DOM-induced hyperthermia but not hypophagia. On the other hand, spiperone (5-HT_{1A}/5-HT_{2A}/D₂ antagonist) pretreatment significantly attenuated DOM-induced hyperthermia but accentuated DOM-induced hypophagia. Daily administration of DOM (1.0 mg kg⁻¹ day⁻¹) produced complete tolerance to its hypophagic effect by day 4 but did not produce cross-tolerance to m-chlorophenylpiperazine-induced hypophagia. In contrast, daily administration of DOM for 7 days did not produce either tolerance to its hyperthermic effect or modify m-chlorophenylpiperazine-induced hyperthermia in rats. These findings suggest that DOM-induced hypophagia and hyperthermia in rats are mediated by stimulation of 5-HT_{2A} receptors.

L8 ANSWER 15 OF 15

MEDLINE on STN

ACCESSION NUMBER:
DOCUMENT NUMBER:
TITLE:

94138685
Pubmed ID: 8306109
Evidence that RU 24969-induced locomotor activity in C57/Bl/6 mice is specifically mediated by the 5-HT_{1B} receptor.

AUTHOR:

Cheetham S C; Heal D J

CORPORATE SOURCE:

Boots Pharmaceuticals Research Department, Nottingham.
British Journal of pharmacology, (1993 Dec) Vol. 110, No. 4, pp. 1621-9.

SOURCE:

Journal code: 7502536. ISSN: 0007-1188.

PUB. COUNTRY:

ENGLAND; United Kingdom

DOCUMENT TYPE:

Journal; Article; (JOURNAL ARTICLE)

LANGUAGE:

English

FILE SEGMENT:

Priority Journals

ENTRY MONTH:

199403

ENTRY DATE:

Entered STN: 30 Mar 1994
Last Updated on STN: 30 Mar 1994
Entered Medline: 17 Mar 1994

AB

1. The behavioural effects of the 5-HT_{1B} receptor agonists, RU 24969 and CGS 12066B, have been investigated in C57/Bl/6 mice. 2. RU 24969 (1-30 mg kg⁻¹) produced intense and prolonged hyperlocomotion and other behavioural changes. 3. CGS 12066B caused similar effects, but they were much less pronounced, inconsistent and transient irrespective of whether this drug was given i.p. (1-15 mg kg⁻¹) or i.c.v. (0.2-40 micrograms). However, CGS 12066B (7.5 and 15 mg kg⁻¹) caused a dose-related inhibition of RU 24969 (7.5 mg kg⁻¹)-induced hyperlocomotion indicating that the former is a 5-HT_{1B} partial agonist. 4. RU 24969 (7.5 mg kg⁻¹ i.p.)-induced hyperlocomotion was inhibited by the (-)-, but not (+)-isomers of pindolol (4 mg kg⁻¹) and propranolol (20 mg kg⁻¹) but not by metoprolol (10 mg kg⁻¹) or ICI 118,551 (5 mg kg⁻¹), consistent with an involvement of 5-HT_{1A} or 5-HT_{1B} receptors. 5. The response was not altered by the selective 5-HT_{1A} receptor antagonist, WAY 100135 (5 mg kg⁻¹, s.c.), the 5-HT_{2A}/5-HT_{2C} receptor antagonist, ritanserin (0.1 mg kg⁻¹), the selective 5-HT₃ receptor antagonist, ondansetron (1 mg kg⁻¹) or the non-selective 5-HT receptor antagonists metergoline (3 mg kg⁻¹) and metergoline (3 mg kg⁻¹). 6. Although spiroxatrine (0.1 mg kg⁻¹) and ketanserin (1 mg kg⁻¹) inhibited RU 24969-induced hyperlocomotion, these effects were probably due to antagonism of dopamine D₂ receptors and alpha 1-adrenoceptors respectively. (ABSTRACT TRUNCATED AT 250 WORDS)

=>

10/800,328

EAST Search History

Ref #	Hits	Search Query	DBs	Default Operator	Plurals	Time Stamp
L1	1	("5157034").PN.	USPAT	OR	OFF	2007/04/09 09:38
L2	877056	514/217.07 OR 514/249 OR 540/599 IR 544/349	US-PGPUB; USPAT; USOCR; FPRS; EPO; JPO; DERWENT	OR	ON	2007/04/09 09:39
L3	✓ 17	L2 AND (PYRIDO[1,2-A]PYRAZINE OR PYRIDO[1,2-A]PYRAZIN)	US-PGPUB; USPAT; USOCR; FPRS; EPO; JPO; DERWENT	OR	ON	2007/04/09 09:40